

STUDY PROTOCOL

Open Access



LOGGIC/FIREFLY-2: a phase 3, randomized trial of tovorafenib vs. chemotherapy in pediatric and young adult patients with newly diagnosed low-grade glioma harboring an activating *RAF* alteration

Cornelis M. van Tilburg^{1,2,3,4,5}, Lindsay B. Kilburn⁶, Sébastien Perreault⁷, Rene Schmidt⁸, Amedeo A. Azizi⁹, Ofelia Cruz-Martínez¹⁰, Michal Zápotočský¹¹, Katrin Scheinemann^{12,13,14}, Antoinette Y. N. Schouten-van Meeteren¹⁵, Astrid Sehested¹⁶, Enrico Opocher¹⁷, Pablo Hernáiz Driever¹⁸, Shivaram Avula¹⁹, David S. Ziegler^{20,21,22}, David Capper^{23,24}, Arend Koch²³, Felix Sahm²⁵, Jiaheng Qiu²⁶, Li-Pen Tsao²⁶, Samuel C. Blackman²⁶, Peter Manley²⁶, Till Milde^{1,2,3,4,5}, Ruth Witt^{1,2,4,5}, David T. W. Jones^{1,4,27}, Darren Hargrave²⁸ and Olaf Witt^{1,2,3,4,5*}

Abstract

Background Pediatric low-grade glioma (pLGG) is essentially a single pathway disease, with most tumors driven by genomic alterations affecting the mitogen-activated protein kinase/ERK (MAPK) pathway, predominantly *KIAA1549::BRAF* fusions and *BRAF* V600E mutations. This makes pLGG an ideal candidate for MAPK pathway-targeted treatments. The type I *BRAF* inhibitor, dabrafenib, in combination with the MEK inhibitor, trametinib, has been approved by the United States Food and Drug Administration for the systemic treatment of *BRAF* V600E-mutated pLGG. However, this combination is not approved for the treatment of patients with tumors harboring *BRAF* fusions as type I *RAF* inhibitors are ineffective in this setting and may paradoxically enhance tumor growth. The type II *RAF* inhibitor, tovorafenib (formerly DAY101, TAK-580, MLN2480), has shown promising activity and good tolerability in patients with *BRAF*-altered pLGG in the phase 2 FIREFLY-1 study, with an objective response rate (ORR) per Response Assessment in Neuro-Oncology high-grade glioma (RANO-HGG) criteria of 67%. Tumor response was independent of histologic subtype, *BRAF* alteration type (fusion vs. mutation), number of prior lines of therapy, and prior MAPK-pathway inhibitor use.

Methods LOGGIC/FIREFLY-2 is a two-arm, randomized, open-label, multicenter, global, phase 3 trial to evaluate the efficacy, safety, and tolerability of tovorafenib monotherapy vs. current standard of care (SoC) chemotherapy in patients < 25 years of age with pLGG harboring an activating *RAF* alteration who require first-line systemic therapy. Patients are randomized 1:1 to either tovorafenib, administered once weekly at 420 mg/m² (not to exceed 600 mg),

*Correspondence:

Olaf Witt

o.witt@kitz-heidelberg.de

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

or investigator's choice of prespecified SoC chemotherapy regimens. The primary objective is to compare ORR between the two treatment arms, as assessed by independent review per RANO-LGG criteria. Secondary objectives include comparisons of progression-free survival, duration of response, safety, neurologic function, and clinical benefit rate.

Discussion The promising tovorafenib activity data, CNS-penetration properties, strong scientific rationale combined with the manageable tolerability and safety profile seen in patients with pLGG led to the SIOPe-BTG-LGG working group to nominate tovorafenib for comparison with SoC chemotherapy in this first-line phase 3 trial. The efficacy, safety, and functional response data generated from the trial may define a new SoC treatment for newly diagnosed pLGG.

Trial registration ClinicalTrials.gov: NCT05566795. Registered on October 4, 2022.

Keywords Chemotherapy, First-line, Pediatric low-grade glioma, pLGG, Tovorafenib, Child, BRAF, MAPK

Background

Pediatric low-grade glioma (pLGG) is a heterogeneous group of World Health Organization (WHO) grade 1 and 2 tumors comprising several subgroups and is the most frequent pediatric central nervous system (CNS) tumor diagnosis, with 1200 to 1500 new cases per year in the United States (US) [1]. Although pLGG is a low-grade tumor with an excellent 10-year overall survival (OS) rate of 94%, the 10-year progression-free survival (PFS) rate in case of an indication for systemic treatment with standard of care (SoC) chemotherapy is only 44% [2, 3]. Therefore, pLGG may be considered a chronic disease in patients whose tumors are unresectable or cannot be completely resected, who will often require several treatment lines throughout life [4]. The late adverse effects of the disease and treatment of it in combination with the damage to important functional structures puts a heavy burden on patients and can lead to loss of visual function and impairment of neurologic, endocrine, and cognitive functioning [5–7].

The current SoC first-line systemic treatment is chemotherapy, most commonly a combination of vincristine and carboplatin (V/C). Although the regimens used by the European Society for Paediatric Oncology—Brain Tumour Group—low-grade glioma working group (SIOPe-BTG LGG WG) and the Children's Oncology Group (COG) differ slightly, both regimens show a similar outcome [8, 9]. The SIOPe-BTG LGG WG reported a response rate of 29% (24 weeks after treatment start) and a five-year PFS rate of 45% [9], whereas the COG reported a response rate of 35% (end of treatment) and a five-year event-free survival rate of 39% [8]. A single-arm phase 2 study of single-agent vinblastine (VBL) resulted in a 19% response rate and a five-year PFS rate of 53% [10]. Several countries/institutions now utilize VBL as first-line therapy on the basis of this trial.

Treatment with V/C or VBL may be accompanied by significant adverse effects, including bone marrow toxicity, neurotoxicity, hearing loss, renal dysfunction, and allergic reactions [3]. The burden of toxicity, frequent hospital visits as well as a significant number of patients progressing after first-line treatment, motivated the search for alternative strategies. In addition, these chemotherapy regimens appear even less effective in infants with pLGG [11], underscoring the need for novel treatment options in this particular subgroup of patients with poor prognosis [9].

Pediatric low-grade glioma is predominantly a single pathway disease driven by alterations in the mitogen-activated protein kinase (MAPK) signaling pathway (also known as the RAS-RAF-MEK-ERK pathway). *KIAA1549::BRAF* fusions, BRAF V600E mutations, *FGFR1* alterations and loss of function mutations of the neurofibromin 1 (*NF1*) gene are the most frequent molecular alterations [12–18]. In addition, oncogene-induced senescence and the senescence-associated secretory phenotype have recently been reported to play an important role in pLGG [19, 20].

As a single pathway disease, pLGG is an ideal candidate for the development of targeted treatments. In a phase 2 trial comparing the combination of the type I BRAF inhibitor dabrafenib and the MEK inhibitor trametinib to chemotherapy in patients with BRAF V600E-mutated pLGG requiring first-line systemic treatment, an objective response rate (ORR) of 47% and a median PFS of 20.1 months were observed, whereas SoC chemotherapy treatment with V/C resulted in an ORR of 11% and a median of PFS 7.4 months [21]. Subsequently, the US Food and Drug Administration (FDA) approved dabrafenib combined with trametinib as a systemic treatment for BRAF V600E-mutated pLGG [22]. The use of type I BRAF inhibitors is limited to patients with tumors harboring a BRAF V600E mutation due to the

risk of paradoxical activation of the MAPK pathway and accelerated tumor growth if used in patients with tumors harboring a *RAF* fusion, such as those involving *BRAF* or *CRAF/RAF1* [23]. In recurrent or progressive disease, MEK inhibitors, such as selumetinib and trametinib, have shown activity [24–26]. However, data generated from ongoing studies of various MAPK pathway inhibitors have shown that responses in pLGGs driven by either *BRAF* V600E mutations (treated with type I *RAF* inhibitors and/or MEK inhibitors) or *KIAA1549::BRAF* fusions (treated with MEK inhibitors) are often only durable for as long as the drug can be administered [27, 28]. Furthermore, while the ORRs of MEK inhibitors are encouraging, responses can be relatively slow [25], and current MEK inhibitors are poorly brain-penetrant and associated with significant peripheral toxicities, mostly dermatological, but also cardiac and ophthalmological adverse events (AEs) [29, 30]. Thus, high target selectivity, CNS penetration properties, short time to response, and favorable tolerability over a long course of treatment are key determinants of sustained clinical activity and therefore treatment success [26].

Tovorafenib (formerly DAY101, TAK-580, MLN2480) is an investigational, oral, brain-penetrant, selective, small molecule, type II *RAF* inhibitor. In contrast to the approved type I *RAF* inhibitors, tovorafenib inhibits both wild-type *BRAF* and *CRAF/RAF1* kinases and, importantly, hyperactivated signaling resulting from *BRAF* fusions, including the *KIAA1549::BRAF* fusion [31].

Tovorafenib was shown to inhibit the kinase activity of *BRAF* kinase domain fusions with various 5' gene partners, most notably fusion with the *KIAA1549* gene. In cellular assays, tovorafenib inhibited *KIAA1549::BRAF* fusion kinase activity with comparable potency to inhibition of *BRAF* V600E and without the paradoxical activation of the MAPK pathway reported for type I *BRAF* inhibitors [32]. Tovorafenib blocked downstream pERK signaling and had less severe dermatological, cardiac, or ophthalmological toxicities compared to other *RAF* or MEK inhibitors [33–35]. Finally, tovorafenib had greater CNS penetration compared with the type I *RAF* inhibitor dabrafenib [32]. The clinical activity of tovorafenib in the currently ongoing phase 2 FIREFLY-1 (PNOC026) study in pediatric patients with *BRAF*-altered, recurrent, or progressive pLGG harboring a *BRAF* fusion or *BRAF* V600E mutation was recently reported. In the registrational arm (69 evaluable patients), the ORR primary endpoint as determined by Response Assessment in Neuro-Oncology high-grade glioma (RANO-HGG) criteria per independent radiology review committee (IRC) was 67% [33, 35]. Responses were observed in patients with tumors harboring *BRAF* V600E mutations who had received prior MAPK-targeted therapy. Finally, the median time to response was 2.8 months according

to RANO-HGG criteria [33, 35], which is shorter in comparison to MEK inhibitors [25] and may have beneficial impact on the functional outcomes of patients.

The promising tovorafenib activity, manageable safety profile, oral availability, once weekly (QW) dosing as well as strong scientific rationale are the basis for the ongoing LOGGIC/FIREFLY-2 trial. The purpose of this trial is to evaluate the efficacy, safety, and tolerability of oral tovorafenib monotherapy versus SoC intravenous chemotherapy in patients with pLGG harboring an activating *RAF* alteration requiring first-line systemic therapy. The primary efficacy endpoint, ORR, will be evaluated for tovorafenib versus SoC chemotherapy as determined by an IRC using RANO-LGG criteria [36]. In addition, PFS will be a key secondary endpoint tested in a hierarchical manner following ORR for the final assessment of efficacy. Importantly, this study also includes endpoints to evaluate improvements in neurologic outcomes, visual function in patients with optic pathway glioma (OPG), and patient-reported outcomes to better describe the overall impact on patients' lives and activities of daily living. Lastly, independent of this trial, tumor material will be submitted to the molecular platform LOGGIC Core BioClinical Data Bank [37], by sites participating in this parallel study, to explore the identification of prognostic and predictive molecular biomarkers including the recently described MAPK inhibitor sensitivity score [38].

Methods/design

Study design

LOGGIC/FIREFLY-2 (NCT05566795) is a two-arm, randomized, open-label, multicenter, global, phase 3 trial to evaluate the efficacy, safety, and tolerability of tovorafenib monotherapy vs. SoC chemotherapy in patients with pLGG harboring an activating *RAF* alteration who require first-line systemic therapy. Approximately 400 treatment-naïve patients will be randomized at a ratio of 1:1 to either Arm 1 (tovorafenib) or Arm 2 (investigator's choice of SoC chemotherapy) (Fig. 1). SoC chemotherapy will consist of either COG-V/C [39], SIOPE-LGG-V/C [9] or VBL [10].

Prior to any study treatment administration, patients fulfilling all eligibility criteria will be centrally randomized to a treatment arm using the IRT (interactive response technology). Randomization between treatment arms will be stratified by primary location of the tumor (supratentorial midline vs. other), type of genomic *RAF* alteration (fusion vs. mutation), *CDKN2A* status (deletion vs. wild-type/unknown), and infant chiasmatic-hypothalamic glioma (CHG) diagnosis (yes vs. no). Patients who experience progressive disease in Arm 2 can crossover to Arm 1 and receive tovorafenib. No allocation concealment will be necessary as this is an open-label study.




Fig. 1 Design of the LOGICC/FIREFLY-2 (NCT05566795) trial. The study consists of a screening phase, a treatment phase, an end of treatment visit, a 30-day safety follow-up visit, and a long-term follow-up period. The total length of the study, from screening through to the end of the study is expected to be seven years. CHG: Chiasmatic-hypothalamic glioma; COG-V/C: Children's Oncology Group-vincristine/carboplatin; LGG: Low-grade glioma; QW: Once weekly; SIOPe-LGG-V/C: Society of Paediatric Oncology Europe-low-grade glioma vincristine/carboplatin

Study objectives

Primary objective

The primary objective is to compare the ORR of tovorafenib monotherapy (Arm 1) vs. investigator's choice of SoC chemotherapy (Arm 2) as assessed by the IRC per RANO-LGG [36] criteria in pediatric and young adult patients with LGG harboring an activating *RAF* alteration who require first-line systemic therapy. The IRC will review all images and response assessments will be determined by the IRC.

Key secondary objectives

- Comparison of PFS between study arms as assessed by the IRC per RANO-LGG criteria
- Comparison of duration of response (DoR) between study arms as assessed by the IRC per RANO-LGG criteria
- Comparison of OS between study arms
- Comparison of safety and tolerability between study arms by assessment of AEs, serious AEs, treatment-emergent AEs, laboratory values, and vital signs

Other secondary objectives

- Comparison of neurologic function in the following domains between study arms: motor function, daily living skills, communication, and socialization using the Vineland Adaptive Behavior Composite Scales (VABS)

- Comparison of changes in visual acuity outcomes between study arms in patients with OPG
- Comparison of ORR between study arms as assessed by the IRC per RANO-HGG [40] criteria and by Response Assessment in Pediatric Neuro-Oncology for LGG (RAPNO-LGG) [41] criteria
- Comparison of clinical benefit rate (CBR) between study arms as assessed by the IRC per RANO-LGG, RANO-HGG, and RAPNO-LGG criteria
- Comparison of time to response (TTR) between study arms as assessed by the IRC per RANO-LGG, RANO-HGG, and RAPNO-LGG criteria
- Comparison of PFS between study arms as assessed by the IRC per RANO-HGG and RAPNO-LGG criteria
- Comparison of DoR between study arms as assessed by the IRC per RANO-HGG and RAPNO-LGG criteria

Exploratory objectives

- Comparison of ORR, CBR, TTR, PFS, and DoR between study arms as assessed by investigators per RANO-LGG criteria
- Comparison of changes in growth and development of patients between study arms
- Comparison of chemotherapy-induced peripheral neuropathy outcomes between study arms by pediatric-modified total neuropathy score in patients \geq five years of age

- Comparison of neuro-endocrine morbidity between study arms
- Comparison of changes in total tumor volume following treatment between study arms
- Comparison of changes in apparent diffusion coefficients within the tumor following treatment between study arms using diffusion-weighted imaging analysis based on MRI scan data
- Patient-reported outcomes (PRO) assessments: comparison of changes in quality of life (QoL) and health utilities measures between study arms using the Pediatrics Quality of Life™-Core Module (PedsQL-Core), Pediatrics Quality of Life™-Cancer (PedsQL-Cancer), and Patient-Reported Outcomes Measurement Information System (PROMIS®) assessment
- Comparison of time to initiation of next treatment following discontinuation of primary therapy between study arms
- Comparison of cystic involution (the change in total tumor volume including possible cystic parts) between study arms measured using MRI and assessed by the IRC
- Comparison of the efficacy and safety of individual SoC chemotherapy regimens vs. tovorafenib
- Determination of the ORR and disease control rate of patients who begin tovorafenib after discontinuing SoC chemotherapy due to radiographic progression as assessed by the IRC per RANO-LGG, RANO-HGG, and RAPNO-LGG criteria, and by the investigators per RANO-LGG criteria
- Evaluation of the concordance of prior local laboratory *RAF* molecular profiling with a central *RAF* alteration assay being evaluated by the sponsor (Day One Biopharmaceuticals)
- Explore whether early responses in infant CHG at six and 12 weeks correlate with response after 24 weeks of treatment
- Assessment of the pharmacokinetics (PK) of tovorafenib through blood sample collection from patients randomized to the tovorafenib arm (Arm 1)

Study population

The phase 3 LOGGIC/FIREFLY-2 in front-line pLGG is enrolling globally and the study is ongoing; the first patient was dosed in March 2023 [42]. The accrual period is estimated to be ~two years to achieve a total of 400 evaluable patients pooled over both arms [43]. The trial is being conducted at academic centers with patients being recruited from ~100 sites, including the SIOPE LOGGIC (Low Grade Glioma in Children) consortium in Europe, the Asia-Pacific region (Singapore, South Korea, Australia and New Zealand), and North America

(Canada and the US). Patients are eligible for inclusion in the study if they are <25 years of age with a histopathological diagnosis of LGG or neuroepithelial tumor (grade 1–2 per the 2021 WHO classification for CNS; [44]), harboring a known activating *RAF* alteration as identified through molecular assays routinely performed at Clinical Laboratory Improvement Amendments (CLIA) of 1988 or other similarly certified laboratories. Patients must be evaluable by RANO-LGG, that is, they must have at least one measurable lesion imaged up to 28 days prior to treatment initiation that can be defined by T2/FLAIR and reproducibly measured by MRI in at least two-dimensions of at least 10 mm in size, and visible on two or more axial slices that are preferably ≤ 5 mm apart with 0 mm skip. Additionally, an indication for first-line systemic therapy is required, meaning that the tumor is non-resectable (either completely unresectable or partially resected with residual tumor that can no longer be resected) at the time of enrollment, and the patient qualifies for one of the following tumor-related indications for first-line drug treatment:

- At primary diagnosis:
 - Present with CHG and be < one year of age at diagnosis, independent of neurologic and/or visual symptoms
 - Diencephalic syndrome
 - Patients with OPG meeting visual-related criteria
 - Neurologic symptoms/deficits
- After completion of an initial observation phase (so called “watch and wait” strategy):
 - Manifestation of diencephalic syndrome
 - Patients with OPG exhibiting progression of visual impairment
 - Deterioration of neurologic symptoms
 - Radiologic progression (local investigator judgement without pre-specification)

Patients < 16 years of age and patients ≥ 16 years of age, will be required to exhibit Lansky and Karnofsky performance status scores, respectively, of $\geq 70\%$. Furthermore, for children and adolescents, their parents/legal guardians must agree to comply with study procedures, including treatment, laboratory monitoring, and required clinic visits for the duration of their participation in the study.

Key exclusion criteria include diagnosis of a patient with any of the following tumor histological types: schwannoma, subependymal giant cell astrocytoma (tuberous sclerosis), or diffuse intrinsic pontine glioma, even if histologically diagnosed as WHO grade 1–2.

Patients with *NF1*-driven tumors are excluded. Furthermore, patients with a tumor exhibiting activating molecular alterations in addition to activating *RAF* alterations are also ineligible for participation even if the tumors are histologically of a low-grade. Activating molecular alterations leading to exclusion include, but are not limited to:

- *IDH1/2* mutations
- Histone H3 mutations
- *NF1* loss-of-function mutations
- *MYBL* alterations
- *FGFR* mutations or fusions
- Diagnosis, or suspected diagnosis of patients with neurofibromatosis type 1 or 2 via genetic testing or current diagnostic clinical criteria would also lead to exclusion.

A full list of inclusion and exclusion criteria are included in Additional File 1.

Arms and interventions

Arm 1

Patients randomized to Arm 1 receive tovorafenib at a dose of 420 mg/m² (not to exceed 600 mg), in the form of a tablet or liquid suspension, QW, continuously on days 1, 8, 15, and 22 of 28-day (four week) cycles. Dosage is adjusted for body surface area (BSA) calculated per the Mösteller Formula [$\sqrt{((\text{height} \times \text{weight})/3600)}$]. Patients with a BSA of 0.6 m² or less will be required to receive the liquid suspension formulation. If appropriate in the opinion of the investigator, patients with a BSA of 0.7 m² or greater may change formulations at any point during treatment. Administration of tovorafenib will continue until radiographic progression based on RANO-LGG criteria as determined by the investigator and confirmed by the IRC, unacceptable toxicity, or the patient withdraws consent (for treatment schema, see Additional File 2). Patients may continue on tovorafenib even following radiographic progression at the investigator's discretion if there is evidence that the patient is still deriving clinical benefit.

Arm 2

Patients randomized to Arm 2 are treated with one of three SoC chemotherapy regimens based on the investigator's choice: COG-V/C regimen, SIOPe-LGG-V/C regimen, or VBL regimen [9, 11, 39]. The SoC chemotherapy regimen administered to each patient is determined prior to randomization (for treatment schema, see Additional File 2). Patients randomized to an Arm 2, receiving the COG-V/C regimen are treated with vincristine (BSA ≥ 0.6 mg/m²: 1.5 mg/m²/day without exceeding 2 mg/day; BSA < 0.6 mg/m²: 0.05 mg/kg/day to 0.80 mg/

day) administered intravenously (IV) QW through weeks 1–10 of a 12-week induction period; and carboplatin (BSA ≥ 0.6 mg/m²: 175 mg/m²/day; BSA < 0.6 mg/m²: 175 mg/m² to 90 mg) administered IV once during weeks 1, 2, 3, 4, 7, 8, 9, and 10. Patients do not receive any treatment for the last two weeks of the induction period. During the maintenance period (eight six-weekly cycles from week 12 to completion of treatment at 60 weeks), vincristine is administered on weeks 1, 2, and 3 and carboplatin on weeks 1, 2, 3, and 4.

Patients randomized to Arm 2 receiving the SIOPe-LGG-V/C regimen undergo an induction period of 24 weeks (7 cycles), during which they receive vincristine (body weight ≥ 10 kg: 1.5 mg/m²/day without exceeding 2 mg/day; body weight < 10 kg: 0.05 mg/kg/day; age < 6 months: further 1/3 dose reduction) IV QW from weeks 1–10, and on weeks 13, 17, and 21. During this period, carboplatin (body weight ≥ 10 kg: 550 mg/m²/day, without exceeding 1050 mg; body weight < 10 kg: 18.3 mg/kg/day; age < 6 months: further 1/3 dose reduction) is administered as a one-hour infusion IV on weeks 1, 4, 7, 10, 13, 17, and 21. Following a four-week period where patients do not receive any treatment, they then enter a consolidation phase until completion of treatment at 81 weeks, during which they receive vincristine and carboplatin in six-week cycles, with vincristine administered on weeks 1, 2, and 3; and carboplatin on week 1.

The Arm 2 VBL regimen comprises VBL (BSA ≥ 0.6 m²: 6 mg/m²/day, without exceeding 10 mg; BSA < 0.6 m²: BSA dose $\times 1/30 \times$ body weight [kg]/day) administered IV QW in four-week cycles until completion of treatment at 70 weeks.

For patients experiencing treatment delays or disruptions, investigators may extend treatment with SoC chemotherapies for up to two additional cycles. Other than the end of study, treatment may be discontinued if a patient exhibits radiographic progression based on RANO-LGG criteria as determined by the investigator and confirmed by the IRC, unacceptable toxicity, or the patient withdraws consent. Patients in Arm 2 who demonstrate radiographic progression during the treatment phase or after completion of chemotherapy may be eligible to receive tovorafenib (Fig. 1).

Dose modifications

If patients in Arm 1 experience an AE that is clinically or medically intolerable based on evaluation by National Cancer Institute Common Terminology Criteria for Adverse Events Version 5.0 (which include AEs that have previously been observed with either tovorafenib administration or other therapies that have a similar mechanism of action), tovorafenib dosing will be interrupted until resolution to grade 1 or the AE reverts to a

baseline level. Upon resolution and/or reversion, dosing may be restarted at the same dose or a lower dose, at the discretion of the local investigator. If the dose is reduced, re-escalation to a higher dose may be permitted after approval by the sponsor's medical monitor or designee. The initial starting dose (420 mg/m² [not to exceed a dose of 600 mg]) is based on BSA. The reconstituted liquid formulation (25 mg/mL) is required for a BSA 0.3 to 0.6 m². For tablets (100 mg) and a BSA 0.9–1.9 m², the first dose reduction is 100 mg from starting dose; the second dose reduction is 100 mg from the first dose reduction (i.e., 200 mg total from the initial starting dose). For the liquid formulation, dose reductions vary from 1 mL (BSAs of 0.3 and 0.5 m²), 2 mL (BSAs of 0.4, 0.6–0.8 m²), 3 mL (BSAs of 0.9–1.2 m²) and 4 mL (BSAs of 1.3–1.9 m²) for the first dose reduction from the starting dose and 1 mL (BSAs of 0.3 and 0.4 m²), 2 mL (BSAs of 0.5–0.9 m²), 3 mL (BSAs of 1.0–1.3 m²) and 4 mL (BSAs of 1.4–1.9 m²) for the second dose reduction from the first dose reduction (i.e., between 2 mL [BSA 0.3 m²] to 8 mL [BSAs of 1.4–1.9 m²] total from the initial starting dose). Chemotherapy dosing for patients in Arm 2 may be reduced or temporarily delayed due to toxicity in accordance with protocol-defined criteria; re-escalation to a higher dose following resolution or reversion of the AE to baseline may be permitted only after approval by the sponsor's medical monitor or designee. All patients on either treatment arm who experience drug-related toxicity requiring a recovery period longer than 42 days will be withdrawn from study drug administration unless, for patients in Arm 1 receiving tovorafenib, there is evidence of clinical benefit, and no alternative treatment is available as determined by the investigator and approved by the sponsor's medical monitor or designee.

Tests and evaluations

Physical evaluations

All radiographic tumor measurements will be carried out using MRI of the brain and/or spine. This will occur up to 28 days prior to starting treatment, and approximately every 12 weeks throughout the treatment and long-term follow-up (LTFU) periods. Neurologic function measured by VABS (motor function, daily living skills, communication and socialization) will be evaluated one, two, and five years after treatment initiation. In addition, screening for visual acuity is required for all patients; baseline assessments will be taken during screening. For patients in Arm 1 (and Arm 2 cross-over patients receiving tovorafenib) with deficiencies in visual function related to OPG, visual acuity testing involving a fundus examination with comment on retinal abnormalities and optic disc, and if possible, visual fields to confrontation, and best corrected visual acuity (BCVA) via logarithm

of the minimum angle of resolution assessment will be performed at every radiographic response assessment, the end-of-treatment visit, and every six months during the LTFU period. For all other patients in Arm 1 and all patients in Arm 2, fundus examinations with a comment on retinal abnormalities and BCVA testing will be completed as needed during patient visits.

In addition to standard ophthalmology examination, monitoring for safety will include physical examination, neurologic examination, dermatology examination, cardiac evaluation, bone assessment (for patients with Tanner stage < 4–5), Karnofsky/Lansky performance score, clinical AEs, laboratory variables and vital signs, including height and weight.

QoL evaluations

PedsQL-Core, PedsQL-Cancer and PROMIS® are being assessed at Screening, every 52 weeks from week 5 to end of treatment, and at year 5 as part of the LTFU. Arm 2 has an additional assessment at year 2 in the LTFU. This difference is because Arm 1 involves treating to progression. All PROs are administered according to the scale developer's intended age group. For children under four years of age, caregiver(s)/parent(s) will complete the proxy-reporting PROs. For children above five years of age, both the child and parent will complete self-report and parent proxy report version of the scale, if applicable. PROs will only be conducted if local language translation is available.

PK evaluations

The dose of tovorafenib will be taken in the clinic on PK sampling days to ensure that PK samples are collected at scheduled times. The schedule of blood sample collections is cycle 1 day 1, cycle 2 day 1 (± three-day window), and cycle 4 day 1 (± three-day window): pre-dose and two hours post-dose, and then every subsequent third cycle through to cycle 13 day 1 (± three-day window): pre-dose, and in the event of toxicity. If a patient receiving tovorafenib experiences an AE that fits the criteria of a severe adverse event (SAE) as determined by the investigator, a blood sample will be collected, when clinically feasible, for measurement of drug concentrations at or around that time.

Statistical analysis

To assess the ORR per the primary endpoint, the planned sample size of approximately 400 patients will provide around 85% power to detect a 15% improvement for the tovorafenib arm at a two-tailed level of significance of 0.05, assuming a 30% ORR in the control arm and a dropout rate of up to 10%. To assess PFS as per the key secondary endpoint, the planned sample size

of approximately 400 patients will provide around 85% power to detect a hazard ratio of 0.67 for PFS at a two-tailed level of significance of 0.05, assuming the median PFS for the tovorafenib arm is 4.5 years versus a median of three years for the control arm. The ORR by RANO-LGG (IRC-assessed) primary analysis is expected to occur ~12 months after the last patient is randomized; a PFS analysis will occur later.

For statistical hypothesis testing of efficacy endpoints, a multiple testing procedure will be applied to control the overall type I error rate at a two-sided alpha level of 5% for the hypotheses testing of the primary (ORR per RANO-LGG criteria) and the key secondary endpoints (PFS per RANO-LGG criteria, DoR per RANO-LGG criteria). Analysis of the primary endpoint ORR as assessed by the IRC per RANO-LGG criteria will be done first at a two-sided alpha level of 5%. If this test meets the pre-specified boundary for statistical significance, then the key secondary endpoints will be tested. It should be noted that the statistical analysis plan for the LOGGIC/FIREFLY-2 trial may be subject to change based on emerging data from FIREFLY-1 trial.

Data monitoring

To ensure a high level of safety monitoring during this trial, an independent Data Monitoring Committee (DMC) has been established and will meet periodically to review safety and efficacy data across the entire tovorafenib program. The purpose of the DMC will be to ensure the ethical conduct of the LOGGIC/FIREFLY-2 study and to protect the safety interests of patients in this study. Based on its review, the DMC will provide the Sponsor with recommendations regarding trial modification, continuation, or termination. Roles and responsibilities of DMC and Sponsor as well as meeting schedule and format of information are set forth in a charter.

Discussion

Patients with pLGG are in need of alternative, effective treatments with less toxicity, deeper and more durable activity and importantly, better neurologic, visual, and patient-reported outcomes. In the registrational arm (Arm 1) of the phase 2 FIREFLY-1 trial evaluating tovorafenib in pediatric patients with pLGG, rapid responses were observed with a median time to response of 2.8 months (RANO-HGG), 5.5 months (RAPNO-LGG [pending adjudication]) and 4.2 months (RANO-LGG), as of a December 22, 2022 data cut-off. Overall response rates of 67% (RANO-HGG, confirmed complete response [cCR] or partial response [PR], includes three unconfirmed partial responses [uPRs]), 51% (RAPNO-LGG, cCR, PR or minor response [MR] [pending adjudication], includes four uPR and

four unconfirmed MR [uMR]), and 49% (RANO-LGG, cCR, PR or MR, includes eight uPR and two uMR) were observed with a clinical benefit rate of 93% (RANO-HGG, cCR, PR or stable disease [SD]), 87% (RAPNO-LGG, cCR, PR, MR or SD [pending adjudication]) and 83% (RANO-LGG, cCR, PR, MD or SD) [33]. No difference in response rate was noted for patients that had previously been treated with MAPK pathway inhibitors. Furthermore, of 136 patients treated in Arms 1 and 2 (safety analysis set), only five discontinued treatment due to AEs (of those, four were treatment related); 39 required dose reductions or treatment interruptions due to treatment-related AEs [33, 35].

The promising phase 2 tovorafenib activity data in combination with the tolerability and safety of this oral monotherapy in patients with pLGG led to the decision of the SIOPe-BTG LGG WG to nominate tovorafenib for a comparison with SoC chemotherapy in the first-line phase 3 trial. This resulted in a collaboration with Day One Biopharmaceuticals who became the sponsor of the LOGGIC/FIREFLY-2 trial. Pediatric oncology drug development is hampered by many hurdles and challenges, especially in pediatric neuro-oncology, and the collaboration between an academic group and a biopharmaceutical company as in the LOGGIC/FIREFLY-2 trial accelerates accessibility of innovative drugs for children with cancer and at the same time provides opportunities for the scientific advancement of the field [45].

In the context of pLGG being a chronic disease, in addition to radiologic responses, functional endpoints such as neurologic outcomes, visual function in OPG, and patient-reported outcomes are of extreme importance and will be assessed throughout this study. As this is a registrational trial, the primary efficacy endpoint in this study is based on RANO-LGG criteria, the same criteria used in registrational study CDRB436G2201 (NCT02684058) of dabrafenib in combination with trametinib in pediatric patients with pLGG harboring a BRAF V600 mutation [36, 46]. Radiologic response will also be measured by RAPNO-LGG criteria [41] as a secondary endpoint; data from this trial will contribute towards the clinical validation of the existing RAPNO working group consensus recommendations.

Although not in the study protocol, tissue can be sent from participating sites for molecular profiling via the LOGGIC Core BioClinical Databank [37]. This will not only confirm the mandatory activating *RAF* alterations but will also allow for exploration and correlation of complex biomarkers based on RNA sequence analysis and clinical outcomes. To this aim, a novel MAPK inhibitor sensitivity score was recently developed to guide biomarker driven future trials, such as those investigating relapse and/or combination therapy [38].

The ongoing LOGGIC/FIREFLY-2 trial will determine how the promising phase 2 activity data in combination with the tolerability and safety of tovorafenib will translate when compared with SoC chemotherapy in the first-line treatment setting [47]. Compared with most currently applied chemotherapy regimens, which require in-clinic intravenous treatment, patients randomized to tovorafenib will receive an oral drug that is taken at home once weekly, that is available in both tablet and a pediatric friendly oral suspension that can be also given via a nasogastric or gastric tube, allowing continuation of daily activities such as attending school. For future patients, the efficacy and safety data generated from this study can potentially address the unmet clinical need in the treatment of pLGG and aims at defining the new SoC treatment for this disease in a global effort.

Abbreviations

AEs	Adverse events
BCVA	Best corrected visual acuity
BSA	Body surface area
CBR	Clinical benefit rate
cCR	Confirmed complete response
CHG	Chiasmatic-hypothalamic glioma
CLIA	Clinical Laboratory Improvement Amendments
CNS	Central nervous system
COG-V/C	Children's Oncology Group-vincristine/carboplatin
DMC	Data monitoring committee
DoR	Duration of response
FLAIR	Fluid attenuated inversion recovery
HGG	High-grade glioma
IRC	Independent radiology review committee
IV	Intravenously
LGG	Low-grade glioma
LOGGIC	Low-Grade Glioma in Children
LTFU	Long-term follow up
MAPK	Mitogen-activated protein kinase
MR	Minor response
MRI	Magnetic resonance imaging
ORR	Objective response rate
OPG	Optic pathway glioma
OS	Overall survival
PFS	Progression-free survival
pLGG	Pediatric low-grade glioma
PRO	Patient-reported outcomes
QW	Once weekly
RANO	Response Assessment in Neuro-Oncology
RAPNO	Response Assessment in Pediatric Neuro-Oncology
SAE	Severe adverse event
SD	Stable disease
SIOPe-BTG-LGG WG	European Society for Paediatric Oncology Brain Tumour Group—low-grade glioma working group
SIOPe-LGG-V/C	Society of Paediatric Oncology Europe-low-grade glioma vincristine/carboplatin
SoC	Standard of care
TTR	Time to response
VABS	Vineland Adaptive Behavior Composite Scales
uMR	Unconfirmed minor response
uPR	Unconfirmed partial response
VBL	Vinblastine
WHO	World Health Organization

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12885-024-11820-x>.

Additional file 1.

Additional file 2.

Acknowledgements

Medical writing support was provided by Matt Jones and Nick Davies, of Cancer Communications and Consultancy Ltd (Macclesfield, UK), with funding from Day One Biopharmaceuticals. D.H is supported by funding from the NIHR Great Ormond Street Hospital Biomedical Research Centre. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR, or the Department of Health. D.S.Z is supported by grants from the National Health and Medical Research Council (Synergy Grant #2019056, and Leadership Grant APP2017898) and Cancer Institute New South Wales Program Grant (TPG2037). The FIREFLY-1 data included in this manuscript were the most recent available when it was submitted for review by the journal; a more recent data cut (June 5, 2023) has been published on November 17, 2023 in Nat Med. (Kilburn, LB, Khuong-Quang, D-A., Hansford, JR, et al. The type II RAF inhibitor tovorafenib in relapsed/refractory pediatric low-grade glioma: the phase 2 FIREFLY-1 trial. Nat Med. 2023. <https://doi.org/10.1038/s41591-023-02668-y>).

Authors' contributions

All authors contributed to the conception and design of the protocol. C.M.v.T, P.M and O.W provided administrative support. All authors contributed to writing the main manuscript text. All authors reviewed the manuscript and provided final approval. All authors are accountable for all aspects of the work outlined in this manuscript.

Funding

This study is funded by Day One Biopharmaceuticals. The Clinical Cooperation Unit Pediatric Oncology, German Cancer Research (DFKZ) in Heidelberg, Germany has received funding from Day One Biopharmaceuticals.

Availability of data and materials

Data sharing is not applicable to this article as no datasets have been generated or analyzed. However, in the future, relevant data supporting the findings of the study will be shared in the context of publication(s). To request individual participant data associated with any Day One Biopharmaceuticals clinical trial, please email clinical@dayonebio.com.

Declarations

Ethics approval and consent to participate

The study was conducted in compliance with ICH Good Clinical Practice guidelines and ethical principles described in the Declaration of Helsinki. The study protocol and all amendments were reviewed by the Independent Ethics Committee (IEC) or Institutional Review Board (IRB) for each participating study center. Study centers as of Sept. 6, 2023 include: Perth Childrens Hospital (Nedlands, Australia), Sydney Children's Hospital (Randwick, Australia), The Royal Children's Hospital – Children's Cancer Centre (Parkville, Australia), Women's and Children's Health Network (North Adelaide, Australia), Medizinische Universität Wien (Innsbruck, Austria), Cliniques Universitaires Saint-Luc (Brussels, Belgium), Universitair Ziekenhuis Gent (Ghent, Belgium), Centre Hospitalier de l'Université Laval et Centre Mère-Enfant Soleil (Québec, Canada), CHU Sainte-Justine (Québec, Canada), The Montreal Children's Hospital (Québec, Canada), Fakultní nemocnice Brno—Dětská, Nemocnice (Brno, Czechia), Motol University Hospital (Prague, Czechia), Rigshospitalet (Copenhagen, Denmark), Tampereen Yliopistollinen Sairaala (Tampere, Finland), Hôpital de la Timone (Marseille, France), Institut Curie (Paris, France), Charité Universitätsmedizin Berlin (Berlin, Germany), Evangelische Klinikum Bethel (EvKB) (Bielefeld, Germany), Universitätsklinikum Hamburg-Eppendorf (Hamburg, Germany), Universitätsklinikum Heidelberg (Heidelberg, Germany), Universitätsklinikum Erlangen (Ulm, Germany), Aghia Sofia General Children's Hospital (Athens, Greece), Athens General Children's Hospital (Athens, Greece), Semmelweis Egyetem Tüztoltó utcai

II. Sz. Gyermekgyógyászati Kliniká (Budapest, Hungary), Prinses Máxima Centrum Kinderoncologie (Utrecht, Netherlands), Starship Paediatric Blood & Cancer Center (Grafton, New Zealand), Seoul National University Hospital (Seoul, South Korea), Severance Hospital, (Seoul, South Korea), SingHealth Group—KK Women's and Children's Hospital (Singapore), Univerzitetni Klinični Center Ljubljana (Ljubljana, Slovenia), Hospital Universitari Vall d'Hebrón (Barcelona, Spain), Hospital Sant Joan de Déu Barcelona (Barcelona, Spain), Hospital Infantil Universitario Niño Jesús (Madrid, Spain), Hospital Universitario Virgen del Rocío (Sevilla, Spain), Hospital Universitari I Politècnic La Fe (Valencia, Spain), Astrid Lindgrens Barnsjukhus Solna (Stockholm, Sweden), Centre Hospitalier Universitaire Vaudois Lausanne (Lausanne, Switzerland), Universitaets—Kinderspital Zürich (Zürich, Switzerland), Ann & Robert H. Lurie Children's Hospital of Chicago, (Illinois, United States), Arnold Palmer Hospital for Children (Florida, United States), Children's Hospital Colorado (Colorado, United States), Children's Hospital Los Angeles (California, United States), Children's National Medical Center (The District of Columbia, United States), Dana-Farber Cancer Institute (Massachusetts, United States), Duke Cancer Institute (North Carolina, United States), New York University Langone Health (New York, United States), Phoenix Children's Hospital (Arizona, United States), Riley Hospital for Children at Indiana University Health (Indiana, United States), St. Louis Children's Hospital (Missouri, United States), Texas Children's Hospital (Texas, United States), University of Michigan—C.S. Mott Children's Hospital (Michigan, United States), University of Rochester (New York, United States), and University of Texas Southwestern Medical Center (Texas, United States). Written, informed consent/pediatric assent will be obtained by the investigator and designated site staff and withdrawal of consent/assent will be available at all times during the study.

Consent for publication

Not applicable.

Competing interests

The authors wish to declare the following competing interests: C.M.v.T. has participated in advisory boards for Novartis, Bayer, and Alexion. L.B.K. has received consulting fees from Blueprint Medicine and contracted research from Novartis, Regeneron Pharmaceuticals, Day One Biopharmaceuticals, Spring Works Therapeutics, Bristol Myers Squibb, and SonALAsense and also is a stock shareholder at Onconova Therapeutics. S.P. has participated in advisory boards for Bayer, Alexion, and Eisai and received research support from Novartis, Bayer, and Roche. A.A.A. has participated in an advisory board for Alexion and received travel and scientific grants from Alexion. M.Z. has participated in an advisory board for AstraZeneca. K.S. has participated in advisory boards for Alexion, Servier, Jazz Pharmaceuticals, Novartis, Roche, Bayer, and NovoNordisk. A.S. has participated in advisory boards for Alexion, AstraZeneca Rare Disease, and Day One Biopharmaceuticals. E.O. has participated in an advisory board for Alexion/AstraZeneca Rare Disease. P.H.D. has participated in an advisory board and consultancy for Alexion. D.S.Z. has received consulting fees from Bayer, AstraZeneca, Accendatech, Novartis, Day One Biopharmaceuticals, FivePhusion, Amgen, Alexion, and Norgine and also received research support from Accendatech. D.C. is a co-founder and shareholder for Heidelberg Epignostix GmbH and receives royalties for sale of BRAF V600E specific antibody VE1. F.S. has received honoraria from Bayer and Illumina and is a co-founder and shareholder for Heidelberg Epignostix GmbH. J.Q., L.-P.T., S.C.B., and P.M. are all employees of Day One Biopharmaceuticals and have received Day One Biopharmaceuticals stock and stock options. T.M. has received research grants from Day One Biopharmaceuticals and Biomed Valley Discovery. D.T.W.J. is a co-founder and shareholder for Heidelberg Epignostix GmbH. D.H. has participated in advisory boards for Alexion/AstraZeneca, Bayer, Bristol Myers Squibb, Celgene, Day One Biopharmaceuticals, Janssen, Novartis, and Roche and also received research grants from Alexion/AstraZeneca and Roche. O.W. has participated in advisory boards for Novartis, Janssen, Roche, Bristol Myers Squibb, and AstraZeneca and also received research grants from Day One Biopharmaceuticals, Biomed Valley Discovery, Bristol Myers Squibb, Syndax, and PreComb. The following authors have no competing interests to declare: R.S., O.C., A. Y. N. SvM, S.A., A.K., and R.W.

Author details

¹Hopp Children's Cancer Center Heidelberg (KITZ), Heidelberg, Germany.

²Clinical Cooperation Unit Pediatric Oncology, German Cancer Research Center (DKFZ), Heidelberg, Germany. ³Department of Pediatric Oncology, Hematology, Immunology and Pulmonology, Heidelberg University Hospital, Heidelberg, Germany. ⁴German Cancer Consortium (DKTK), Heidelberg, Germany. ⁵National Center for Tumor Diseases (NCT), Heidelberg, Germany.

⁶Children's National Hospital, Washington, DC, USA. ⁷CHU Sainte-Justine, Université de Montréal, Montréal, QC, Canada. ⁸Institute of Biostatistics

and Clinical Research, Münster, Germany. ⁹Department of Pediatrics and Adolescent Medicine, Comprehensive Center for Pediatrics and Comprehensive Cancer Center, Medical University of Vienna, Vienna, Austria. ¹⁰Neuro-oncology Unit, Pediatric Cancer Center, Hospital Sant Joan de Déu, Barcelona, Spain.

¹¹Department of Paediatric Haematology and Oncology, Charles University, Second Faculty of Medicine and University Hospital Motol, Prague, Czech Republic. ¹²Division of Oncology-Hematology, Children's Hospital of Eastern Switzerland, St. Gallen, Switzerland. ¹³Faculty of Health Sciences and Medicine, University of Lucerne, Lucerne, Switzerland. ¹⁴Department of Pediatrics, McMaster Children's Hospital and McMaster University, Hamilton, Canada.

¹⁵Department of Neuro-oncology, Princess Máxima Center for Pediatric Oncology, Utrecht, The Netherlands. ¹⁶Department of Pediatrics and Adolescent Medicine, Rigshospitalet, Copenhagen, Denmark. ¹⁷Pediatric Hematology, Oncology and Stem Cell Transplant Division, Padua University Hospital, Padua, Italy.

¹⁸German HIT-LOGGIC-Registry for LGG in Children and Adolescents, Charité-Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany. ¹⁹Department of Radiology, Alder Hey Children's Hospital NHS Foundation Trust, Liverpool, UK. ²⁰Kids Cancer Centre, Sydney Children's Hospital, Randwick, NSW, Australia.

²¹Lowy Cancer Research Centre, Children's Cancer Institute, University of New South Wales, Sydney, NSW, Australia. ²²School of Clinical Medicine, University of New South Wales, Sydney, NSW, Australia. ²³Department of Neuropathology, Charité - Universitätsmedizin Berlin, Berlin, Germany. ²⁴DKTK Partner Site, Berlin, Germany. ²⁵Department of Neuropathology, German Cancer Research Center (DKFZ), University Hospital Heidelberg and CCU Neuropathology, German Consortium for Translational Cancer Research (DKTK), Heidelberg, Germany. ²⁶Day One Biopharmaceuticals, Brisbane, CA, USA. ²⁷Division of Pediatric Glioma Research, German Cancer Research Center (DKFZ), Heidelberg, Germany. ²⁸UCL Great Ormond Street Institute of Child Health and Great Ormond Street Hospital for Children, London, UK.

Received: 31 August 2023 Accepted: 2 January 2024

Published online: 30 January 2024

References

- Ostrom QT, Price M, Ryan K, Edelson J, Neff C, Cioffi G, et al. CBTRUS Statistical Report: Pediatric Brain Tumor Foundation Childhood and Adolescent Primary Brain and Other Central Nervous System Tumors Diagnosed in the United States in 2014–2018. *Neuro Oncol*. 2022;24(Suppl 3):iii1–38.
- Gnekow AK, Falkenstein F, von Hornstein S, Zwiener I, Berkefeld S, Bison B, et al. Long-term follow-up of the multicenter, multidisciplinary treatment study HIT-LGG-1996 for low-grade glioma in children and adolescents of the German Speaking Society of Pediatric Oncology and Hematology. *Neuro Oncol*. 2012;14(10):1265–84.
- Gnekow AK, Kandels D, Tilburg CV, Azizi AA, Opocher E, Stokland T, et al. SIOP-E-BTG and GPOH guidelines for diagnosis and treatment of children and adolescents with low grade glioma. *Klin Padiatr*. 2019;231(3):107–35.
- Goebel AM, Gnekow AK, Kandels D, Witt O, Schmidt R, Hernaiz Driever P. Natural history of pediatric low-grade glioma disease - first multi-state model analysis. *J Cancer*. 2019;10(25):6314–26.
- Aleman M, Velasco R, Simo M, Bruna J. Late effects of cancer treatment: consequences for long-term brain cancer survivors. *Neurooncol Pract*. 2021;8(1):18–30.
- Heitzer AM, Ashford JM, Hastings C, Liu APY, Wu S, Bass JK, et al. Neuropsychological outcomes of patients with low-grade glioma diagnosed during the first year of life. *J Neurooncol*. 2019;141(2):413–20.
- Liu APY, Hastings C, Wu S, Bass JK, Heitzer AM, Ashford J, et al. Treatment burden and long-term health deficits of patients with low-grade gliomas or glioneuronal tumors diagnosed during the first year of life. *Cancer*. 2019;125(7):1163–75.
- Ater JL, Zhou T, Holmes E, Mazewski CM, Booth TN, Freyer DR, et al. Randomized study of two chemotherapy regimens for treatment of low-grade glioma in young children: a report from the Children's Oncology Group. *J Clin Oncol*. 2012;30(21):2641–7.
- Gnekow AK, Walker DA, Kandels D, Picton S, Giorgio P, Grill J, et al. A European randomised controlled trial of the addition of etoposide to standard vincristine and carboplatin induction as part of an 18-month treatment programme for childhood (<=16 years) low grade glioma - A final report. *Eur J Cancer*. 2017;81:206–25.

10. Lassaletta A, Scheinemann K, Zelcer SM, Hukin J, Wilson BA, Jabado N, et al. Phase II weekly vinblastine for chemotherapy-naïve children with progressive low-grade glioma: A Canadian Pediatric Brain Tumor Consortium study. *J Clin Oncol*. 2016;34(29):3537–43.
11. Kandels D, Pietsch T, Bison B, Warmuth-Metz M, Thomale UW, Kortmann RD, et al. Loss of efficacy of subsequent nonsurgical therapy after primary treatment failure in pediatric low-grade glioma patients-Report from the German SIO-P-LGG 2004 cohort. *Int J Cancer*. 2020;147(12):3471–89.
12. Fisher MJ, Jones DTW, Li Y, Guo X, Sonawane PS, Waanders AJ, et al. Integrated molecular and clinical analysis of low-grade gliomas in children with neurofibromatosis type 1 (NF1). *Acta Neuropathol*. 2021;141(4):605–17.
13. Jones DT, Hutter B, Jager N, Korshunov A, Kool M, Warnatz HJ, et al. Recurrent somatic alterations of FGFR1 and NTRK2 in pilocytic astrocytoma. *Nat Genet*. 2013;45(8):927–32.
14. Jones DT, Kocalkowski S, Liu L, Pearson DM, Backlund LM, Ichimura K, et al. Tandem duplication producing a novel oncogenic BRAF fusion gene defines the majority of pilocytic astrocytomas. *Cancer Res*. 2008;68(21):8673–7.
15. Pfister S, Janzarik WG, Remke M, Ernst A, Werft W, Becker N, et al. BRAF gene duplication constitutes a mechanism of MAPK pathway activation in low-grade astrocytomas. *J Clin Invest*. 2008;118(5):1739–49.
16. Ryall S, Zapotocky M, Fukuoka K, Nobre L, Guerreiro Stucklin A, Bennett J, et al. Integrated molecular and clinical analysis of 1,000 pediatric low-grade gliomas. *Cancer Cell*. 2020;37(4):569–583e565.
17. Sturm D, Pfister SM, Jones DTW. Pediatric gliomas: current concepts on diagnosis, biology, and clinical management. *J Clin Oncol*. 2017;35(21):2370–7.
18. Zhang J, Wu G, Miller CP, Tatevosian RG, Dalton JD, Tang B, et al. Whole-genome sequencing identifies genetic alterations in pediatric low-grade gliomas. *Nat Genet*. 2013;45(6):602–12.
19. Buhl JL, Selt F, Hielscher T, Guiho R, Ecker J, Sahm F, et al. The senescence-associated secretory phenotype mediates oncogene-induced senescence in pediatric pilocytic astrocytoma. *Clin Cancer Res*. 2019;25(6):1851–66.
20. Selt F, Sigaud R, Valinciute G, Sievers P, Zaman J, Alcon C, et al. BH3 mimetics targeting BCL-XL impact the senescent compartment of pilocytic astrocytoma. *Neuro Oncol*. 2023;25(4):735–47.
21. Bouffet E, Hansford J, Garré ML, Hara J, Plant-Fox A, Aerts I, et al. Primary analysis of a phase II trial of dabrafenib plus trametinib (dab + tram) in BRAF V600-mutant pediatric low-grade glioma (pLGG). *J Clin Oncol*. 2022;40(Suppl 17):LBA2002–LBA2002.
22. Tafinlar (dabrafenib) prescribing information. https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/202806s022lbl.pdf. Accessed 4 June 2023.
23. Schreck KC, Grossman SA, Pratilas CA: BRAF mutations and the utility of RAF and MEK inhibitors in primary brain tumors. *Cancers (Basel)*. 2019;11(9).
24. Bouffet E. Trametinib therapy in pediatric patients with low-grade gliomas (LGG) with BRAF gene fusion; a disease-specific cohort in the first pediatric testing of trametinib. *Neuro Oncol*. 2018;20(Suppl 2):i114.
25. Fangusaro J, Onar-Thomas A, Young Poussaint T, Wu S, Ligon AH, Lindeman N, et al. Selumetinib in paediatric patients with BRAF-aberrant or neurofibromatosis type 1-associated recurrent, refractory, or progressive low-grade glioma: a multicentre, phase 2 trial. *Lancet Oncol*. 2019;20(7):1011–22.
26. Selt F, van Tilburg CM, Bison B, Sievers P, Harting I, Ecker J, et al. Response to trametinib treatment in progressive pediatric low-grade glioma patients. *J Neurooncol*. 2020;149(3):499–510.
27. Rosenberg T, Yeo KK, Mauguén A, Alexandrescu S, Prabhu SP, Tsai JW, et al. Upfront molecular targeted therapy for the treatment of BRAF-mutant pediatric high-grade glioma. *Neuro Oncol*. 2022;24(11):1964–75.
28. Stege H, Haist M, Schultheis M, Fleischer MI, Mohr P, Meier F et al: Discontinuation of BRAF/MEK-directed targeted therapy after complete remission of metastatic melanoma-a retrospective multicenter ADOReg study. *Cancers (Basel)*. 2021;13(10).
29. Klesse LJ, Jordan JT, Radtke HB, Rosser T, Schorry E, Ullrich N, et al. The use of MEK inhibitors in Neurofibromatosis type 1-associated tumors and management of toxicities. *Oncologist*. 2020;25(7):e1109–16.
30. Lugowska I, Kosela-Paterczyk H, Kozak K, Rutkowski P. Trametinib: a MEK inhibitor for management of metastatic melanoma. *Onco Targets Ther*. 2015;8:2251–9.
31. Olszanski AJ, Gonzalez R, Corrie P, Pavlick AC, Middleton M, Lorigan P et al: Phase I study of the investigational, oral pan-RAF kinase inhibitor TAK580 (MLN2480) in patients with advanced solid tumors (ST) or melanoma (MEL): Final analysis. *Ann Oncol*. 2017;28.
32. Sun Y, Alberta JA, Pilarz C, Calligaris D, Chadwick EJ, Ramkissoon SH, et al. A brain-penetrant RAF dimer antagonist for the noncanonical BRAF oncoprotein of pediatric low-grade astrocytomas. *Neuro Oncol*. 2017;19(6):774–85.
33. Kilburn L, Khuong-Quang D-A, Nysom K, Landi D, Ziegler D, Hernáiz Driever P et al: Clinical activity of pan-RAF inhibitor tovorafenib in the registrational pediatric low-grade glioma arm of the phase 2 FIREFLY-1 (PNOC026) study. *J Clin Oncol*. 2023;41 Suppl 16:abstr 10004 and associated presentation.
34. Rasco DW, Medina T, Corrie P, Pavlick AC, Middleton MR, Lorigan P, et al. Phase 1 study of the pan-RAF inhibitor tovorafenib in patients with advanced solid tumors followed by dose expansion in patients with metastatic melanoma. *Cancer Chemother Pharmacol*. 2023;92(1):15–28.
35. Kilburn L, Khuong-Quang D-A, Nysom K, Landi D, Ziegler D, Hernáiz Driever P et al: Clinical activity of pan-RAF inhibitor tovorafenib in the registrational pediatric low-grade glioma arm of the phase 2 FIREFLY-1 (PNOC026) study. *Neuro Oncol*. 2023;25 i57 and associated presentation.
36. van den Bent MJ, Wefel JS, Schiff D, Taphoorn MJ, Jaecle K, Junck L, et al. Response assessment in neuro-oncology (a report of the RANO group): assessment of outcome in trials of diffuse low-grade gliomas. *Lancet Oncol*. 2011;12(6):583–93.
37. Hardin EC, Schmid S, Sommerkamp A, Boddén C, Heipertz AE, Sievers P et al: LOGGIC Core BioClinical Data Bank: Added clinical value of RNA-seq in an international molecular diagnostic registry for pediatric low-grade glioma patients. *Neuro Oncol*. 2023; noad078.
38. Sigaud R, Albert TK, Hess C, Hielscher T, Winkler N, Kocher D, et al. MAPK inhibitor sensitivity scores predict sensitivity driven by the immune infiltration in pediatric low-grade gliomas. *Nat Commun*. 2023;14(1):4533.
39. Packer RJ, Lange B, Ater J, Nicholson HS, Allen J, Walker R, et al. Carboplatin and vincristine for recurrent and newly diagnosed low-grade gliomas of childhood. *J Clin Oncol*. 1993;11(5):850–6.
40. Wen PY, Chang SM, Van den Bent MJ, Vogelbaum MA, Macdonald DR, Lee EQ. Response assessment in neuro-oncology clinical trials. *J Clin Oncol*. 2017;35(21):2439–49.
41. Fangusaro J, Witt O, Hernáiz Driever P, Bag AK, de Blank P, Kadom N, et al. Response assessment in paediatric low-grade glioma: recommendations from the Response Assessment in Pediatric Neuro-Oncology (RAPNO) working group. *Lancet Oncol*. 2020;21(6):e305–16.
42. Day One Bio March 6, 2023 press release. <https://ir.dayonebio.com/news-releases/news-release-details/day-one-reports-fourth-quarter-and-full-year-2022-financial>. Accessed 14 December 2023.
43. DAY101 vs. Standard of Care Chemotherapy in Pediatric Patients With Low-Grade Glioma Requiring First-Line Systemic Therapy (LOGGIC/FIREFLY-2). <https://clinicaltrials.gov/ct2/show/NCT05566795>. Accessed 4 June 2023.
44. Louis DN, Perry A, Wesseling P, Brat DJ, Cree IA, Figarella-Branger D, et al. The 2021 WHO Classification of Tumors of the Central Nervous System: a summary. *Neuro Oncol*. 2021;23(8):1231–51.
45. Vassal G, de Rojas T, Pearson ADJ. Impact of the EU Paediatric Medicine Regulation on new anti-cancer medicines for the treatment of children and adolescents. *Lancet Child Adolesc Health*. 2023;7(3):214–22.
46. Dabrafenib With Trametinib for Pediatric Low-Grade Glioma With BRAF V600E Mutation. <https://ascopost.com/issues/april-25-2023/dabrafenib-with-trametinib-for-pediatric-low-grade-glioma-with-braf-v600e-mutation/>. Accessed 16 August 2023.
47. Kilburn L, Landi D, Leary S, Ziegler D, Baxter P, Franson A, et al. FIREFLY-1 (PNOC026): Phase 2 study of pan-Raf inhibitor tovorafenib in pediatric and young adult patients with Raf-altered recurrent or progressive low-grade glioma or advanced solid tumors. *Neuro Oncol*. 2022;24:89.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.