

SPECIAL REPORT

Post-diagnosis physical activity and sedentary behaviour and colorectal cancer prognosis: A Global Cancer Update Programme (CUP Global) systematic literature review and meta-analysis

Georgios Markozannes^{1,2}  | Nerea Becerra-Tomás¹ | Margarita Cariolou¹ |
 Katia Balducci¹ | Rita Vieira¹ | Sonia Kiss¹ | Dagfinn Aune^{1,3,4}  |
 Darren C. Greenwood⁵ | Marc J. Gunter^{1,6} | Ellen Copson⁷ |
 Andrew G. Renehan⁸ | Martijn Bours⁹ | Wendy Demark-Wahnefried¹⁰  |
 Melissa M. Hudson¹¹ | Anne M. May¹² | Folakemi T. Odedina¹³ |
 Roderick Skinner¹⁴ | Karen Steindorf¹⁵ | Anne Tjønneland^{16,17} |
 Galina Velikova¹⁸ | Monica L. Baskin¹⁹ | Rajiv Chowdhury²⁰ | Lynette Hill²¹ |
 Sarah J. Lewis²² | Jaap Seidell²³ | Matty P. Weijenberg⁹ | John Krebs²⁴ |
 Amanda J. Cross¹ | Konstantinos K. Tsilidis^{1,2}  | Doris S. M. Chan¹ 

Correspondence

Doris S. M. Chan, Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, St. Mary's Campus, Norfolk Place, London, W2 1PG, UK.
 Email: d.chan@imperial.ac.uk

Funding information

World Cancer Research Fund; Wereld Kanker Onderzoek Fonds; American Institute for Cancer Research

Abstract

Low physical activity and high sedentary behaviour have been clearly linked with colorectal cancer development, yet data on their potential role in colorectal cancer survival is limited. Better characterisation of these relationships is needed for the development of post-diagnosis physical activity and sedentary behaviour guidance for colorectal cancer survivors. We searched PubMed and Embase through 28 February 2022 for studies assessing post-diagnosis physical activity, and/or sedentary behaviour in relation to all-cause and cause-specific mortality and recurrence after colorectal cancer diagnosis. Total and recreational physical activity were assessed overall and by frequency, duration, intensity, and volume using categorical, linear, and non-linear dose-response random-effects meta-analyses. The Global Cancer Update Programme (CUP Global) independent Expert Committee on Cancer Survivorship and Expert Panel interpreted and graded the likelihood of causality. We identified 16 observational studies on 82,220 non-overlapping patients from six countries. Physical activity was consistently inversely associated

Konstantinos K. Tsilidis and Doris S. M. Chan contributed equally and shared last authorship.

Where authors are identified as personnel of the International Agency for Research on Cancer/World Health Organization, the authors alone are responsible for the views expressed in this article and they do not necessarily represent the decisions, policy or views of the International Agency for Research on Cancer/World Health Organization.

For affiliations refer to page 438

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Authors. *International Journal of Cancer* published by John Wiley & Sons Ltd on behalf of UICC.

with colorectal cancer morbidity and mortality outcomes, with 13%–60% estimated reductions in risk. Sedentary behaviour was positively associated with all-cause mortality. The evidence had methodological limitations including potential confounding, selection bias and reverse causation, coupled with a limited number of studies for most associations. The CUP Global Expert panel concluded limited-suggestive evidence for recreational physical activity with all-cause mortality and cancer recurrence. Total physical activity and its specific domains and dimensions, and sedentary behaviour were all graded as limited-no conclusion for all outcomes. Future research should focus on randomised trials, while observational studies should obtain objective and repeated physical activity measures and better adjustment for confounders.

KEYWORDS

colorectal cancer survival, evidence grading, meta-analysis, physical activity, systematic review

What's new?

The roles of physical activity and sedentary behaviour in colorectal cancer prognosis are unclear. Here, as part of CUP Global, the authors performed a systematic review and meta-analysis on the associations of post-diagnosis physical activity and sedentary behaviour with all-cause and cause-specific mortality and recurrence among colorectal cancer survivors. Whilst the evidence was graded as limited, it suggested that a physically active lifestyle and avoidance of sedentary behaviour may be associated with longer overall survival. The study highlights the importance of more consistent and objective exposure assessment in future studies and the need for randomised trials to provide stronger evidence.

1 | INTRODUCTION

Colorectal cancer is the third leading cause of cancer incidence and the second leading cause of cancer-related mortality worldwide, with an estimated >1.9 million incident cases and >0.9 million deaths, respectively, in 2020.¹ The effectiveness of early detection via cancer screening programmes in reducing mortality^{2–4} coupled with better cancer treatment options have led to increased survival rates among individuals diagnosed with colorectal cancer.⁵ The estimated 5-year survival for colorectal cancer is generally over 60% in western populations and high income countries,^{5–7} and on average 20% lower in transitioning countries.⁸ Among patients diagnosed with earlier stages of the disease, survival rates are even higher, and the 5-year prevalence of the disease is estimated to be over 5.2 million people worldwide, making colorectal cancer the second most prevalent cancer, after breast cancer.¹

Extensive research has been conducted to identify risk factors affecting the incidence of colorectal cancer. In addition to genetic predisposition and increasing age, several environmental and lifestyle factors have been associated with the development of the disease. Among these, the protective role of physical activity on colorectal cancer risk, specifically on colon cancer, is well-established.^{9,10} The Third Expert Report from the World Cancer Research Fund/American Institute for Cancer Research (WCRF/AICR) concluded that there is convincing evidence that physical activity reduces colon cancer

incidence,⁹ while the 2018 Physical Activity Guidelines Advisory Committee (PAGAC) Scientific Report¹¹ graded this association as being supported by strong evidence.

Despite the abundance of evidence on colorectal cancer incidence, evidence regarding the association between physical activity and sedentary behaviour and colorectal cancer prognosis is relatively limited. A small number of meta-analyses investigating the role of post-diagnosis physical activity on outcomes after colorectal cancer^{12–18} suggested an inverse association with all-cause mortality. These meta-analyses have primarily focused on the association of categories of physical activity levels with colorectal cancer prognosis, but none explicitly assessed the different dimensions and domains of activity. Moreover, apart from one ongoing trial¹⁹ there is a lack of randomised controlled trials (RCTs) assessing the effect of physical activity on colorectal cancer survival. The role of sedentary behaviour as a distinct risk factor from physical activity, defined as the energy expenditure of less than 1.5 metabolic equivalents of task (METs) while one assumes a sitting or reclining posture,²⁰ is even less studied.²¹

Due to the lack of strong evidence, physical activity and sedentary behaviour recommendations targeting cancer outcomes tailored to cancer survivors are limited. Formal evaluation of the existing observational evidence base performed by PAGAC concluded moderate evidence strength for all-cause and cancer-specific mortality in colorectal cancer, based on individual studies published until

February 2018.¹¹ The 2018 American College of Sports Medicine (ACSM) Roundtable also concluded that physical activity is beneficial for survival²² and other cancer-related health outcomes.²³ However, the precise quantity of physical activity required to effectively reduce cancer-specific or all-cause mortality remains undetermined.²² The evidence on sitting time as a proxy of total sedentary time and cancer-specific mortality was limited.²² Currently, colorectal cancer patients and survivors are advised to follow the same recommendations as for cancer prevention.^{9,24–26}

The aim of WCRF International's new Global Cancer Update Programme (CUP Global) formerly known as the WCRF/AICR Continuous Update Project²⁷ is the development of disease- and stage-specific guidance and recommendations. Therefore, as part of CUP Global, we present the findings of a systematic literature review on the association of post-diagnosis physical activity and sedentary behaviour and colorectal cancer prognosis. Evidence on adiposity and diet and outcomes after colorectal cancer, along with an overall summary are presented in the accompanied papers.^{28–30}

2 | METHODS

This systematic review is part of the ongoing CUP Global programme and is reported following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines³¹ (Supplementary Table S1). The protocol can be found elsewhere.³²

2.1 | Search strategy, selection criteria and data extraction

Comprehensive details on the methodology are available in the Supplementary Texts 1 and 2. Briefly, we searched PubMed and Embase through 28th February 2022. We included observational studies, RCTs, or pooled analyses of such studies evaluating post-diagnosis (at or shortly after diagnosis, during treatment, after the end of treatment) physical activity, sedentary behaviour, or their pre-to-post diagnosis or post-diagnosis changes. We included studies with at least 100 participants with primary colorectal cancer or its subtypes diagnosed during adulthood. The term 'colorectal' referred to any colorectal cancer survivor, regardless of cancer subtype, colon or rectum. Outcomes of interest included all-cause, colorectal cancer site and subsite-specific, and cardiovascular-specific mortality, second primary cancer, and cancer recurrence/disease-free survival. Study selection was performed at title and abstract level, followed by full-text review, and supplemented with manual reference screening. If more than one publication from the same population was identified, we included the one with the higher number of events per outcome assessed. Data were extracted into the CUP Global database and included, among others, publication details, study characteristics (number of participants, events, and inclusion and exclusion criteria), exposure and outcome details, relative risks (RR) 95% confidence intervals (CI),

standard errors, and/or p-values, and details on adjustment variables. Study selection and data extraction were performed by one reviewer and at least 10% checked by another; disagreements were resolved by consensus.

2.2 | Evidence synthesis

We reviewed total physical activity (defined as any combined activity domain, including occupational, recreational, household, transportation) and the specific domains, irrespective of which physical activity dimension was assessed (frequency, duration, intensity). Specifically for recreational physical activity, the most investigated domain, we further reviewed frequency, duration, and intensity (activity of any intensity and specifically moderate-to-vigorous intensity). We also investigated recreational physical activity volume as a separate dimension, measured in MET-h/week. Furthermore, we performed meta-analyses on sedentary behaviour when its definition was sufficiently consistent across studies (capturing similar types of sedentary behaviours).

We analysed cancer recurrence/disease-free survival as defined by the studies (Supplementary Table S2). When more than one of these outcomes were reported, the one with the highest number of events was included in the meta-analyses.

We performed DerSimonian-Laird random-effects meta-analyses³³ for categorical (highest vs. lowest levels) and linear dose-response comparisons when at least two and three studies provided relevant information, respectively. When at least five studies provided sufficient information, we also performed one-stage non-linear dose-response meta-analyses³⁴ using restricted cubic splines with knots placed at the 10th, 50th, and 90th percentiles of the exposure distribution. The nonlinear component was tested using a likelihood ratio test.³⁵

We assessed the presence of small-study effects (such as publication bias) by visual inspection of the funnel-plots and used the Egger's regression asymmetry test,³⁶ when at least 10 studies were included in a meta-analysis. If small study effects bias was suggested, we estimated an adjusted summary RR using the trim-and-fill method.³⁷ Heterogeneity was described by the between-study variation (τ^2) and by the range of study estimates in each meta-analysis. The I^2 metric of inconsistency³⁸ with cut-offs at $I^2 = 30\%$, 50% , and 75% indicated low, moderate, high, and substantial proportion of total variability in effect estimates due to between-study heterogeneity. The 95% prediction intervals depicted the range of estimates likely to contain the value of a new study.³⁹

We performed pre-defined stratified meta-analyses by sex, cancer anatomical subsite and stage, and molecular subtype (based on sequencing or immunohistochemistry approaches). We performed sensitivity meta-analyses by excluding, when possible, survivors with locally advanced and metastatic tumours, to explore potential impact of bias by reverse causation. We also performed leave-one-out sensitivity meta-analyses to assess the magnitude of the effect of exclusion of each study on the summary estimate.⁴⁰

2.3 | Risk of bias assessment

A modified version of the Risk of Bias for Nutrition Observational Studies (RoB-NObs) tool⁴¹ was utilised to assess the risk of bias in the studies included in the meta-analyses. The tool was originally developed by the U.S. Department of Agriculture (USDA) Nutrition Evidence Systematic Review after modifications to the Cochrane's collaboration Risk Of Bias In Non-randomized Studies of Interventions (ROBINS-I).⁴² The tool underwent further refinement and testing by the Imperial College London (ICL) review team to ensure its suitability for investigating exposure-outcome associations in cancer survivorship studies. This involved adapting the tool's prompting questions and providing additional guidance to encompass adiposity, physical activity, and dietary/nutritional exposures (the working document version dated 11/07/2023 can be found in Supplementary Table S3). The tool consists of seven domains, including confounding, participant selection, exposure classification, departures from intended exposures, missing data, outcome measurement, and selective reporting. In the context of cancer survivorship, we have specifically designated age, adiposity, and cancer stage and treatment as pivotal confounding variables a-priori of the studied associations, due to their substantial impact on both the levels of physical activity and sedentary behaviour engagement along with their recognised role as robust prognostic factors for survival in colorectal cancer patients.

2.4 | Evidence grading criteria

The CUP Global independent Expert Committee on Cancer Survivorship and Expert Panel convened by WCRF International interpreted the findings, independently of ICL. The Expert Committee made the preliminary conclusions, and the Expert Panel made the final conclusions. The quality of the evidence was graded for all exposures as strong (subgrades evaluating likelihood of causality: convincing, probable, substantial effect on risk unlikely) or limited (subgrades evaluating likelihood of causality: limited-suggestive, limited-no conclusion) according to a list of pre-defined criteria assessing the quantity, consistency, magnitude, and precision of the summary estimates, evidence of biological gradient, the study design, risk of bias, generalisability, and biological plausibility of the results (Supplementary Table S4). The grades of the quality of the evidence reflect the independent Expert Committee's and Expert Panel's confidence that the association estimates are correct.

3 | RESULTS

A flowchart of the selection process is presented in Figure 1. We excluded 453 publications on exposures other than physical activity, 64 on pre-diagnosis-only exposures,^{43–106} seven on a mixture of pre- and post-diagnosis exposures,^{107–113} and one where the outcome was colorectal polyps.¹¹⁴ Overall, 28 publications met the eligibility criteria. Five publications^{115–118} that were superseded by more recent publications and three^{119–121} only reporting unadjusted results were excluded

from any subsequent analyses. Therefore, a total of 20 publications (16 studies) on post-diagnosis physical activity, pre-to post and post-diagnosis change in physical activity levels, and post-diagnosis sedentary behaviour were included. These pertained to 16 publications of 13 cohort studies^{122–137} and four publications of three observational follow-up analyses of patients enrolled in clinical treatment RCTs^{138–141} not aiming to evaluate physical activity interventions. The number of publications may differ from the number of studies as several publications from the same study reporting on different exposure-outcome associations were included,^{126,131,134–136,139,140} but also one publication included two studies.¹²³ No primary RCT was identified.

Detailed information on study and participants' characteristics and physical activity definitions and domains are presented in Supplementary Tables S5 and S6. The 16 studies comprised 82,220 non-overlapping patients, of whom about 7800 died of any cause, about 1700 died of colorectal cancer, and about 2100 experienced an additional colorectal cancer event. The median study size was 1640 (range 247–43,596) participants in cohort studies and 1218 (832–1992) participants in secondary analyses of RCTs. Most studies ($n = 9$; 60%) were from the United States,^{124,126,128–130,132,135,136,138–141} three were from the Netherlands,^{123,137} and one each from Australia,¹³³ Germany,¹²⁷ Japan,¹²⁵ and South Korea.¹²² Most studies included individuals with stage I–III ($n = 7$ ^{123,126,128,129,131–136}) or I–IV disease ($n = 4$ ^{122,124,127,130}), of which 2 studies reported the percentage of stage IV patients (3.5% and 17%) and another two did not report the %. In addition, two studies included only stage III,^{140,141} one included stage 0–IV,¹²⁵ and two included only survivors with tumour metastases.^{137,138}

Six (38%) studies^{123,127,133,137,141} measured physical activity of any type (total, including multiple types such as recreational, occupational, household, and other types of activity), and 12 (75%) studies^{122,124–126,128,130–132,134–141} measured recreational physical activity. The results of the two studies that further measured other specific types of activity, such as cycling,¹²⁷ gardening,¹²⁷ housework,¹²⁷ sports,¹²⁷ or walking^{127,138} were descriptively synthesised. Eight studies (50%)^{126,127,131,132,134–139,141} assessed total (low, moderate, and vigorous) intensity of physical activity, seven (44%)^{122–124,128,130,133} assessed physical activity of moderate-to-vigorous intensity, while the intensity of physical activity was unclear in one study.¹²⁵ Physical activity was self-reported in all studies but only nine (56%) used validated questionnaires.^{123,126,127,131–138} Two studies^{133,136} examined physical activity changes (pre-to-post-diagnosis or post-diagnosis). Four studies^{127–130} assessed sedentary behaviour as daily time spent sitting while watching TV^{127–129} or as overall sitting time/day.¹³⁰

Figure 2 and Supplementary Table S7 show a summary of the results of the meta-analyses.

3.1 | Physical activity

3.1.1 | Total physical activity

Six studies (five publications) were identified.^{123,127,133,137,141} The category with the highest versus lowest levels of total physical activity was inversely associated with all-cause mortality (RR = 0.70; 95% CI = 0.55–0.89; $I^2 = 47\%$; $\tau^2 = 0.03$; RRs range = 0.48–0.80; six




FIGURE 1 Flowchart of study selection process.

studies; 1443 deaths),^{123,127,133,141} but the evidence for cancer recurrence/disease-free survival was weak ($RR = 0.87$; 95% $CI = 0.63-1.20$; $I^2 = 0\%$; RRs range = $0.75-0.99$; two studies; 762 deaths)^{123,141} (Supplementary Figure S1). Dose-response meta-analysis could not be conducted due to insufficient data.

When excluding survivors with locally advanced and metastatic tumours, the summary RR for all-cause mortality was similar but with lower heterogeneity than when including all colorectal cancer patients ($RR = 0.70$; 95% $CI = 0.56-0.87$; $I^2 = 21\%$; $\tau^2 = 0.09$; RRs range = $0.53-0.80$; four studies, three publications)^{123,127,133}




FIGURE 2 Summary risk ratios (RRs) and 95% confidence intervals (95% CI) for post-diagnosis physical activity and sedentary behaviour and outcomes after colorectal cancer. CI, confidence interval; MET, metabolic equivalent of task; RR, relative risk.

(Supplementary Figure S2). The leave-one-out sensitivity analyses did not identify any influential studies (Supplementary Figure S3).

3.1.2 | Recreational physical activity

Twelve studies (12 publications) were identified.^{122,124,125,128,130,132,135-139,141} Most studies investigated volume of recreational activity (MET-h/week).^{125,130,132,135,136,138-140} Four studies measured duration of recreational activity^{124,128,137,138} and three measured physical activity frequency.^{122,125,141} The primary analysis on recreational physical activity included all studies, irrespective of the dimension of activity measured, however, due to heterogeneity across definitions and dimensions measured, meta-analyses are also presented by specific dimensions.

A lower risk of all-cause mortality was observed when comparing the highest with the lowest levels of recreational physical activity (any dimension) (RR = 0.67; 95% CI = 0.60–0.75; $I^2 = 56\%$; $\tau^2 = 0.02$; RRs range = 0.39–0.85; 12 studies; 9539 deaths)^{122,124,125,128,130,132,135-139,141} (Supplementary Figure S4). We found evidence of small-study effects from visual inspection of the funnel plot and the Egger's test ($p = .002$) (Supplementary Figure S5). The trim and fill sensitivity analysis suggested that after adjusting for six potentially missing studies, the summary RR would be RR = 0.76 (95% CI = 0.68–0.86) (Supplementary Figure S6).

The highest versus lowest levels of recreational physical activity (any dimension) were associated with a lower risk of colorectal cancer-specific mortality (RR = 0.63; 95% CI = 0.47–0.84; $I^2 = 58\%$; $\tau^2 = 0.06$; RRs range = 0.29–0.83; six studies; 2406 deaths)^{122,128,130,132,135,136} and cancer recurrence (RR = 0.80; 95% CI = 0.70–0.92; $I^2 = 14\%$; $\tau^2 = 0.002$; RRs range = 0.52–0.83; three studies, all secondary analyses of RCTs; 4095 events).^{138,140,141} (Supplementary Figure S4).

Since studies often included patients with various cancer stages, the predefined subgroup meta-analyses by stage could not be performed. When excluding survivors with locally advanced and metastatic tumours, the summary RR and heterogeneity for all-cause mortality were similar to the results including all colorectal cancer patients (RR = 0.65; 95% CI = 0.56–0.75; $\tau^2 = 0.02$; RRs range = 0.41–0.78; $I^2 = 54\%$; eight studies)^{122,124,125,128,130,132,135,136} (Supplementary Figure S7). The leave-one-out sensitivity analyses did not identify any influential studies (Supplementary Figure S8).

3.1.3 | Recreational physical activity frequency

Three studies (three publications) were identified.^{122,125,141} The highest versus lowest levels of physical activity frequency showed an inverse association with all-cause mortality (RR = 0.77; 95%

CI = 0.72–0.83; $I^2 = 0\%$; $\tau^2 = 0.00$; RRs range = 0.75–0.78; three studies; 3247 deaths)^{122,125,141} (Supplementary Figure S9). Dose-response meta-analyses were not possible due to limited data.

In subgroup meta-analyses by cancer subsite, the highest versus lowest levels of recreational physical activity frequency were inversely associated with all-cause mortality risk in both colon (RR = 0.77; 95% CI = 0.71–0.84; $I^2 = 0\%$; $\tau^2 = 0.00$; RRs range = 0.66–0.79; three studies; 2502 deaths) and rectal (RR = 0.76; 95% CI = 0.67–0.86; RRs range = 0.75–0.86; $I^2 = 0\%$; two studies; 1135 deaths)^{122,125} cancer survivors (Supplementary Figure S10).

3.1.4 | Recreational physical activity duration

Four studies (four publications) were identified.^{124,128,137,138} Highest versus lowest recreational physical activity duration (minutes or h/week) was inversely associated with the risk of all-cause mortality (RR = 0.66; 95% CI = 0.52–0.84; $I^2 = 51\%$; $\tau^2 = 0.03$; RRs range = 0.39–0.80; three studies; 1820 deaths)^{124,128,137,138} (Supplementary Figure S11). Dose-response meta-analysis was not possible due to insufficient data.

3.1.5 | Recreational physical activity intensity

Six studies (six publications) assessed total recreational physical activity intensity.^{132,135,136,138,139,141} Highest versus lowest levels of recreational activity intensity (total) were inversely associated with all-cause (RR = 0.65; 95% CI = 0.53–0.80; $I^2 = 60\%$; $\tau^2 = 0.03$; RRs range = 0.41–0.85; six studies; 2325 deaths)^{132,135,136,138,139,141} and colorectal cancer-specific mortality (RR = 0.40; 95% CI = 0.25–0.62; $I^2 = 0\%$; $\tau^2 = 0.00$; RRs range = 0.29–0.47; three studies; 211 deaths)^{132,135,136} and cancer recurrence (RR = 0.83; 95% CI = 0.73–0.94; $I^2 = 0\%$; RRs range = 0.82–0.83; two studies; 1744 events)^{138,141} (Supplementary Figure S12).

When excluding survivors with locally advanced and metastatic tumours, the summary RR for all-cause mortality was 0.51 (95% CI = 0.39–0.67) with no heterogeneity ($I^2 = 0\%$; $\tau^2 = 0.00$; RRs range = 0.41–0.59; three studies)^{132,135,136} (Supplementary Figure S13).

Five studies (five publications) assessed moderate-to-vigorous intensity of recreational physical activity.^{122,124,128,130,137} Highest versus lowest levels of recreational physical activity (moderate-to-vigorous) were associated with lower risk of all-cause (RR = 0.48; 95% CI = 0.53–0.78; $I^2 = 68\%$; $\tau^2 = 0.03$; RRs range = 0.39–0.78; four studies; 2445 deaths)^{122,124,128,130,137} and colorectal cancer-specific mortality (RR = 0.82; 95% CI = 0.74–0.89; $I^2 = 0\%$; $\tau^2 = 0.00$; RRs range = 0.53–0.82; three studies; 1040 deaths).^{122,128,130} There was weak (and likely underpowered) evidence of an association with cardiovascular disease-specific mortality (RR = 0.63; 95% CI = 0.33–1.22; $I^2 = 79\%$; $\tau^2 = 0.26$; RRs range = 0.36–0.89; three studies; 287 events)^{122,128,130} (Supplementary Figure S14).

3.1.6 | Recreational physical activity volume

Six studies (seven publications) were identified.^{130,132,135,136,138–140} Highest compared to lowest volumes of recreational physical activity were associated with decreased risk of all-cause (RR = 0.60; 95% CI = 0.48–0.75; $I^2 = 65\%$; $\tau^2 = 0.04$; RRs range = 0.41–0.85; six studies; 2632 deaths)^{130,132,135,136,138,139} and colorectal cancer-specific mortality (RR = 0.52; 95% CI = 0.30–0.87; $I^2 = 62\%$; $\tau^2 = 0.17$; RRs range = 0.29–0.87; four studies; 624 deaths).^{130,132,135,136} The summary RR for high versus low volumes of recreational physical activity and cancer recurrence was 0.73 (95% CI = 0.50–1.06; $I^2 = 56\%$; RRs range = 0.55–0.83; two studies; 1317 events)^{136,138} (Supplementary Figure S15).

In the sensitivity analysis excluding survivors with locally advanced and metastatic tumours, the summary RR for all-cause mortality was similar but with no heterogeneity (RR = 0.55; 95% CI = 0.47–0.65; $I^2 = 0\%$; $\tau^2 = 0.00$; RRs range = 0.41–0.59; four studies)^{130,132,135,136} (Supplementary Figure S16).

In the linear dose-response meta-analyses, an increase of 10 MET-h/week of recreational physical activity was associated with decreased risk of all-cause (RR = 0.76; 95% CI = 0.66–0.88; $I^2 = 86\%$; $\tau^2 = 0.02$; RRs range = 0.53–0.94; six studies; 2632 deaths)^{130,132,135,136,138,139} and colorectal cancer-specific mortality (RR = 0.73; 95% CI = 0.63–0.84; $I^2 = 0\%$; $\tau^2 = 0.00$; RRs range = 0.63–0.84; four studies; 624 deaths)^{130,132,135,136} (Supplementary Figure S17). Linear dose-response meta-analysis was not possible for the other outcomes due to limited data.

In the sensitivity analysis excluding survivors with locally advanced and metastatic tumours, the results for all-cause mortality were similar to the analysis including all colorectal cancer patients (RR = 0.68; 95% CI = 0.55–0.84; $I^2 = 76\%$; $\tau^2 = 0.03$; RRs range = 0.53–0.83; four studies) (Supplementary Figure S18). Sequential omission of each study did not change the results (Supplementary Figure S19).

Non-linear dose-response analysis was only possible for all-cause mortality. The evidence from six studies was suggestive of a non-linear inverse association ($P_{\text{non-linearity}} = .07$; 2408 deaths)^{130,132,135,136,138,139} indicating a consistent inverse association up to 20 MET-h/week and remained relatively constant for higher activity levels (Supplementary Figure S20). There was no strong evidence of non-linearity ($P_{\text{non-linearity}} = .27$; four studies; 1075 deaths) when excluding survivors with locally advanced and metastatic tumours, but the data were sparse^{130,132,135,136} (Supplementary Figure S21).

Subgroup analyses by sex were only possible for all-cause mortality, with high versus low volumes of recreational physical activity showing inverse associations for both men (RR = 0.65; 95% CI = 0.52–0.80; $I^2 = 0\%$; RRs range = 0.59–0.68; two studies; 540 deaths)^{130,135} and women (RR = 0.55; 95% CI = 0.34–0.88; $I^2 = 57\%$; $\tau^2 = 0.10$; RRs range = 0.41–0.81; three studies; 411 deaths)^{130,132,136} (Supplementary Figure S22).

Stratified analyses by molecular subtypes were only performed in subsets of participants ($n = 371$ –605) of the Nurses' Health Study & Health Professionals Follow-Up Study,^{126,131,134} so we descriptively

summarise these results. In general, most molecular subtypes did not show evidence of interaction, except for *IRS1*, *CTNNB1*, *PTGS2*, and *P27* and only for colorectal cancer-specific mortality but not for overall survival (Supplementary Figures S23 and S24).

3.1.7 | Other types of physical activity

A pattern of inverse associations with all-cause mortality was observed across the different types of physical activities analysed (walking, bicycling, gardening, housework, and sport) (RRs ranged from 0.34 to 0.90, with the 95% CI not including the null in two out of the six comparisons)^{127,138} (Supplementary Figure S25). No evidence of association was observed in the only study analysing walking and cancer recurrence.¹³⁸

3.2 | Physical activity change

Two studies (two publications)^{133,136} were identified, which investigated changes in physical activity levels around the time of diagnosis. A meta-analysis was not performed due to large methodological differences, but we descriptively synthesised these results. In one study,¹³³ increases of total physical activity by >0–2 h/week and by ≥2 h/week from pre-diagnosis to 5 months post-diagnosis compared to no change or decreased levels showed no strong evidence of association with all-cause mortality (RR = 1.27; 95% CI = 0.88–1.83 and RR = 1.06; 95% CI = 0.65–1.71, respectively). There was some weak evidence that a post-diagnosis increase (from 5 to 12 months) by 0–2 h/week was associated with a lower risk of all-cause mortality (RR = 0.79; 95% CI = 0.59–1.04), while an increase by ≥2 h/week was associated with a lower risk of all-cause mortality (RR = 0.69; 95% CI = 0.50–0.94) compared to no change or decreased activity levels. In the Nurses' Health Study,¹³⁶ an increase of at least one level of recreational physical activity (defined as categories of MET-h/week) from a median of 6 months before to a median of 22 months post-diagnosis compared to no change in activity levels over this period was associated with lower all-cause mortality risk (RR = 0.53; 95% CI = 0.30–0.85), while a decrease of physical activity by one level compared to no change showed no association (RR = 1.23; 95% CI = 0.79–1.91).

3.3 | Sedentary behaviour

Four studies (four publications) of self-reported sedentary behaviour were identified.^{127–130} We performed a meta-analysis of the three that estimated time sitting while watching TV. Highest versus lowest levels of time sitting while watching TV were associated with 28% higher risk of all-cause mortality (RR = 1.28; 95% CI = 1.06–1.55; I^2 = 0%; τ^2 = 0.00; RRs range = 1.16–1.45; three studies; 969 deaths)^{127–129} and showed imprecise evidence of a higher risk of colorectal cancer-specific mortality (RR = 1.45; 95%

CI = 0.95–2.21; I^2 = 0%; RRs range = 1.45–1.45; two studies; 252 deaths)^{128,129} (Supplementary Figure S26). Linear dose-response meta-analysis was only possible for all-cause mortality. An increase of 120 min/day of sitting while watching TV was associated with higher risk of all-cause mortality (RR = 1.13; 95% CI = 1.02–1.26; I^2 = 0%; τ^2 = 0.00; RRs range = 1.09–1.40; three studies; 969 deaths) (Supplementary Figure S27).

3.4 | Risk of bias

Around 37% of the publications on physical activity had moderate RoB in confounding, 21% had serious, and around 42% had critical RoB by failing to adjust for important confounders (i.e., age, stage, cancer treatment). Participation in all the studies was conditional on survival and self-selection cannot be excluded, therefore, all publications had a serious RoB in this domain, except for one that utilised data from a national registry thus accounting for selection bias.¹²² Most publications (74%) had a moderate RoB in exposure misclassification, mainly due to the very detailed information provided on exposure assessment, despite many utilising non-validated tools. All publications had a critical RoB in departures from intended exposures as time-varying analyses were not performed, except for one that partially accounted for deviations of the intended measurements.¹³⁹ Most publications (58%) had a serious RoB in missing data and 21% had a low RoB either due to no missingness or because appropriate statistical methods were utilised to account for missingness. Most publications (84%) had a low RoB in outcome measurement; the rest did not provide sufficient information for assessment. All publications had a moderate RoB in selective reporting (Supplementary Figures S28 and S29).

For sedentary behaviour, 75% had a critical RoB in confounding by not adjusting for all important confounders, all had serious RoB in participant selection, all had either serious or critical RoB in exposure misclassification, all had critical RoB in departures from the intended exposures, all but one¹²⁸ which accounted for missingness had a serious RoB due to missing data, all had low RoB in outcome measurement and all had moderate RoB in selective reporting (Supplementary Figures S28 and S29).

3.5 | Evidence grading

Table 1 reports the evidence grading. Relatively sparse data, methodological and clinical considerations, inconsistencies in the assessment of physical activity, increased likelihood of residual confounding from comorbidities and other factors, and potential for reverse causation were among the reasons that led to limited conclusions on the likelihood of causality for the existing evidence. The evidence on post-diagnosis recreational physical activity and all-cause mortality and cancer recurrence received the limited-suggestive sub-grade. A limited-no conclusion sub-grade was given to recreational physical activity and other outcomes apart from all-cause mortality and cancer

TABLE 1 Summary of the judgement of the Independent WCRF Expert Panel.

Physical activity and sedentary behaviour and survival in colorectal cancer						
		Exposure	Exposure contrast	Outcomes	Summary of findings	Conclusions
					N studies, N events; RR (95% CI); I ² (%); tau ² ; Egger's p-value ^a	
Strong evidence	Convincing	–	–	–	–	–
	Probable	–	–	–	–	–
Limited evidence	Limited—suggestive	Decreases risk				
		Recreational physical activity				
		Any definition	Highest vs. lowest	All-cause mortality	12 studies, 9539 deaths, 0.67 (0.60, 0.75), I ² = 56%, tau ² = 0.02, P _{Egger} < 0.01	The evidence is modest, substantial heterogeneity was observed along with evidence of publication bias, and possible methodological issues.
				Cancer recurrence	3 studies, 4095 events, 0.80 (0.70, 0.92), I ² = 14%, tau ² = 0.002	The evidence rises from three secondary analyses of randomized controlled trials which were relatively consistent in terms of populations (advanced and metastatic cases) and timing of exposure measurement and with low heterogeneity.
	Limited—no conclusion	Total physical activity				The evidence for total physical activity and recreational physical activity dimensions is sparse, often with substantial heterogeneity, and is generally limited by methodological issues including heterogeneity in exposure definition and timing of measurement, and quality issues including increased risk of survival and reverse causation bias.
		Any definition	Highest vs. lowest	All-cause mortality	6 studies, 1443 deaths, 0.70 (0.55, 0.89), I ² = 47%, tau ² = 0.03	
				Cancer recurrence	3 studies, 762 events, 0.87 (0.63, 1.20), I ² = 0%, tau ² = NA	
		Recreational physical activity				
		Any definition	Highest vs. lowest	Colorectal cancer mortality	6 studies, 2406 deaths, 0.63 (0.47, 0.84), I ² = 58%, tau ² = 0.06	
		Frequency	Highest vs. lowest	All-cause mortality	3 studies, 3247 deaths, 0.77 (0.72, 0.83), I ² = 0%, tau ² = 0.00	
		Duration	Highest vs. lowest	All-cause mortality	4 studies, 1920 deaths, 0.66 (0.52, 0.84), I ² = 51%, tau ² = 0.03	
		Intensity (any)	Highest vs. lowest	All-cause mortality	6 studies, 2325 deaths, 0.65 (0.53, 0.80), I ² = 60%, tau ² = 0.03	
				Colorectal cancer mortality	3 studies, 211 deaths, 0.40 (0.25, 0.62), I ² = 0%, tau ² = 0.00	
				Cancer recurrence	2 studies, 1744 events, 0.83 (0.73, 0.94), I ² = 0%, tau ² = 0.00	
		Intensity (moderate-to-vigorous)	Highest vs. lowest	All-cause mortality	5 studies, 2445 deaths, 0.64 (0.53, 0.78), I ² = 68%, tau ² = 0.03	

TABLE 1 (Continued)

Physical activity and sedentary behaviour and survival in colorectal cancer				
Exposure	Exposure contrast	Outcomes	Summary of findings	Conclusions
			N studies, N events; RR (95% CI); I^2 (%); τ^2 ; Egger's p -value ^a	
Volume		Colorectal cancer mortality	3 studies, 1040 deaths, 0.82 (0.74, 0.89), I^2 = 0%, τ^2 = 0.00	
		CVD mortality	3 studies, 287 deaths, 0.63 (0.33, 1.22), I^2 = 79%, τ^2 = 0.26	
	Highest vs. lowest	All-cause mortality	6 studies, 2632 deaths, 0.60 (0.48, 0.75), I^2 = 65%, τ^2 = 0.04	
	10 MET-hours/week		6 studies, 2632 deaths, 0.76 (0.66, 0.88), I^2 = 86%, τ^2 = 0.02	
	Highest vs. lowest	Colorectal cancer mortality	4 studies, 624 deaths, 0.52 (0.30, 0.87), I^2 = 62%, τ^2 = 0.17	
	10 MET-hours/week		4 studies, 624 deaths, 0.73 (0.63, 0.84), I^2 = 0%, τ^2 = 0.00	
Physical activity change	Highest vs. lowest	Cancer recurrence	2 studies, 1317 events, 0.73 (0.50, 1.06), I^2 = 55%, τ^2 = NA	The evidence for changes in physical activity is weak, heterogeneous, and inconsistent, and likely affected by biases.
	Pre-to-post diagnosis	All-cause mortality	3 RRs from 2 studies for moderate and high increases vs no change, no meta-analysis. 2 RRs ranging 1.06 to 1.27, both CIs crossing the null & 1 RR=0.53, CI not crossing the null.	
	Post-diagnosis	All-cause mortality	2 RRs from 1 study for moderate and high increases vs no change, no meta-analysis. 2 RRs ranging 0.69 to 0.79, 1 CI crossing the null.	
Sedentary behaviour				The body of evidence is weak showing higher risk of all-cause mortality and indicating a higher risk for colorectal cancer-specific mortality with higher levels of sedentary behaviour, which may be influenced by reverse causation and other biases.
Sitting while watching TV	Highest vs. lowest	All-cause mortality	3 studies, 969 deaths, 1.28 (1.06, 1.55), I^2 = 0%, τ^2 = 0.00	
	120 min/week		3 studies, 969 deaths, 1.13 (1.02, 1.26), I^2 = 0%, τ^2 = 0.00	
	Highest vs. lowest	Colorectal cancer mortality	2 studies, 252 deaths, 1.45 (0.95, 2.21), I^2 = 0%, τ^2 = NA	

Abbreviations: CI, confidence interval; CVD, cardiovascular disease; NA, not applicable; RR, relative risk.

^aEstimated when at least 10 studies were available.

recurrence, and to recreational physical activity dimensions (volume, frequency, intensity, duration), total physical activity, change of physical activity, and sedentary behaviour.

4 | DISCUSSION

This systematic literature review showed that higher levels of recreational physical activity and its different dimensions were consistently inversely associated with major clinical outcomes in colorectal cancer patients. Due to methodological limitations leading to increased likelihood of biases, small number of studies, and often high between-study variability due to heterogeneity, the CUP Global independent Expert Committee and Expert Panel concluded that the evidence on post-colorectal cancer diagnosis physical activity and sedentary behaviour and survival-related outcomes was limited. The association of recreational physical activity with all-cause mortality that was supported by a relatively larger number of studies and without high levels of between-study heterogeneity, along with the evidence on cancer recurrence that was supported by relatively more consistent studies in terms of population characteristics and timing of exposure measurement reached a limited-suggestive conclusion for the likelihood on causality.

The evidence on recreational physical activity volume also suggested an inverse non-linear relationship with all-cause mortality up to 20 MET-h/week (roughly equivalent to 5 h of moderate-intensity physical activity/week¹⁴²) compared to zero MET-h/week, which plateaued for higher volumes, although no evidence of non-linearity was found in a sensitivity analysis excluding survivors with locally advanced and metastatic tumours. There was less available evidence for total activity, dimensions of recreational activity, change in activity, and sedentary behaviour on survival-related outcomes and no conclusion could be reached. The evidence base was limited and did not allow for comprehensive evaluation of most of the a-priori defined subgroups analyses.

A plethora of mechanisms have been proposed to explain the link between physical activity and colorectal carcinogenesis,¹⁴³ such as changes in the gastrointestinal transit time, immune function, bile acid secretion, lipid and hormone profiles, and insulin and insulin-like growth factor (IGF) alterations. Some of these mechanisms including changes to the IGF-axis and immune response have also been hypothesised to affect colorectal cancer survival, albeit supported by less data,^{144–146} while new proposed pathways of action include exercise-induced shear stress, extracellular vesicles, and systemic milieu alterations.¹⁴⁴ Increased blood pressure and heart rate during exercise leads to higher blood flow and shear pressure,¹⁴⁷ which may reduce the number of circulating tumour cells by immediate cell necrosis promotion.^{148,149} The beneficial role of physical activity in body composition and its anti-inflammatory effects via enhancement of skeletal muscle has also been proposed.^{150–152} For example, exercise leads to increased levels of serum adiponectin and decreased levels of insulin and leptin, which may lead to reduced proliferation and enhanced apoptosis of tumour cells, although evidence

specifically in colorectal cancer patients is currently limited.¹⁵³ Extracellular vesicles, small membrane-surrounded structures which transfer molecules across cells, are an emerging target for treatment of various cancers.^{154,155} Physical activity enhances skeletal muscle mass, which increases extracellular vesicle secretion in blood, leading to increased circulation of beneficial myokines, cytokines and peptides released from muscle cells, such as interleukin-6 (IL-6). Exercise-induced myokine secretion has been shown to have an anti-inflammatory role in various chronic diseases.¹⁵⁶ Physical activity also promotes immune cell mobilisation, which, coupled with increased exercise-induced IL-6 circulation, may lead to increased immune cell tumour infiltration and enhance anti-cancer immunity.¹⁵⁷ Finally, IGF-I and -II and IGF binding protein 3 (IGFBP-3) have been linked with colorectal tumour severity.^{158–160} Despite the inconsistent response of IGFs to physical activity in healthy individuals, limited evidence supports the beneficial role of exercise on colorectal cancer mortality via IGF pathway alterations.^{65,66} Nevertheless, the diverse mechanisms through which physical activity is linked to outcomes after cancer may be influenced by the interplay between activity and treatment in cancer survivors, while the existence of comorbidities may also affect the observed effects through similar biological pathways.

To the best of our knowledge, our analysis is the most comprehensive evaluation of the existing body of evidence on post-diagnosis physical activity and sedentary behaviour and colorectal cancer outcomes. Our results are in line with the findings of previous reviews and meta-analyses of physical activity in the field^{12–18} in both direction and magnitude. However, in the previous meta-analyses, studies assessing total and recreational physical activities were frequently combined. The most recent analysis pooling studies of both post-diagnosis total and recreational physical activity found a 37% and a 21% reduced risk of all-cause mortality for highest versus lowest levels (10 studies, any dimension) and for 10 MET-h/week increase (seven studies),¹⁸ respectively. Those results were similar in magnitude to ours on recreational physical activity volume (40% and 24%, respectively). Half to two-thirds of the studies within these meta-analyses employed validated instruments for the quantification of physical activity, reducing the probability of exposure misclassification and consequently enhancing the reliability of the findings. Change in physical activity has been previously meta-analysed using the same two identified studies.¹⁵ In the present work, due to the high clinical and methodological heterogeneity across studies, including largely different definitions and timeframes of assessment, we decided against a quantitative synthesis. We identified one meta-analysis on post-diagnosis sedentary behaviour and colorectal cancer-specific mortality showing similar results to ours, however sitting while watching TV and overall sitting time were analysed together.²¹

Apart from the updated evidence base, our review is the first to assess specific physical activity domains and dimensions in relation to colorectal cancer outcomes to increase homogeneity of exposure assessment, as well as to evaluate non-linearity. Furthermore, the totality of the evidence was evaluated by the CUP Global independent Expert Committee and Expert Panel following a rigorous evaluation using pre-defined standardised grading criteria. Nevertheless, the

results presented should be interpreted considering certain limitations. Cancer survivorship studies of observational nature are subject to a range of methodological limitations, survivor bias for instance, which could not be excluded in most studies. In addition, there was an increased RoB due to incomplete confounding adjustment in most studies. Studies often did not adjust for important confounders such as treatment and/or stage, factors which can have a large effect on both the exposures and outcomes of interest, while most studies did not account for the presence of comorbidities, which may have an important impact on the ability to participate in physical activity and survival. Finally, the literature search ended on 28 February 2022. Thereby, any relevant studies published after this date were not included. Nonetheless, given the pivotal role RCTs in our evidence grading framework, we performed a comprehensive literature review targeting RCTs published subsequently and until August 31, 2023. However, we did not identify any published physical activity trial among colorectal cancer survivors, apart from the currently ongoing CHALLENGE trial. That trial has an estimated completion date in late 2030.^{19,161} As such, we anticipate that the conclusions would remain unaltered.

There are complexities in determining the feasibility of trials for survival-related outcomes. Deciding whether mortality should be the primary focus is not a straightforward decision. To conduct intention-to-treat analyses, participants should ideally remain actively engaged until reaching the primary endpoint. Additionally, selecting and maintaining the appropriate level of activity for the comparison group is challenging, particularly considering existing recommendations for physical activity for cancer survivors and the potential for contamination between groups. The evidence, however, is supported by studies of different designs including clinical treatment trials with a secondary aim of evaluating the impact of physical activity on cancer outcomes. The populations of these studies were more homogeneous, composed of individuals with advanced and metastatic tumours and the timing of exposure measurement was also more consistent, capturing activity close to treatment.

Physical activity was not consistently assessed across observational studies, as they used different tools for measuring activity levels, measured different types of activity, and assessed different aspects. Our review explicitly assessed several different aspects of physical activity, including frequency, intensity, volume, and duration to enhance definition homogeneity, but studies did not mutually adjust for any of these factors. All studies utilised self-report questionnaires, of which only around half have been validated. Although all studies provided comprehensive descriptions of exposure assessment, physical activity is an umbrella term that describes a breadth of complex behaviours, often inhibiting direct comparability across studies. Even within the same domain or dimension, the comprehensiveness of the assessment varied across studies. Furthermore, self-reported physical activity and sedentary behaviour have been shown to have poor agreement with accelerometer-based measurements generally,¹⁶² and specifically in colon cancer patients,¹⁶³ leading to an increased likelihood of exposure misclassification. This misclassification is expected to be non-differential which may result in attenuated

associations. The wide range of tools used for exposure assessment and the inclusion of cancer patients of any stage may partly contribute to the heterogeneity of the results across studies. Finally, while TV viewing time was used as a surrogate measure of sedentary behaviour, the two are not directly equivalent, since other sedentariness aspects such as occupational sedentary time are not captured in TV viewing time.

The literature is primarily focused on the assessment of recreational activity. Evidence on total activity is sparse while there is some limited evidence on other types of activity, such as household activities. The timing of physical activity measurement differed across studies, ranging from immediately after diagnosis to over 10 years post-diagnosis. Individuals are more likely to engage in lower levels of activity during and short-term after cancer treatment,^{164,165} and their attitudes towards exercise engagement during this period are mixed.^{166,167} In contrast, studies on long-term survivors, that usually exclude patients with relapse or with most aggressive or treatment-resistant disease, indicate good proportions of adherence to physical activity recommendations.^{168,169} Such inconsistencies across the timing of measurement may contribute to the observed heterogeneity and aggravate direct comparisons across studies but due to the large differences and low number of studies, subgroup analyses were not feasible. Moreover, almost all studies used a single measurement as the exposure. Colorectal cancer patients have inconsistent levels of activity throughout the course of the disease, and it has been suggested that even if patients engage in a short-term physical activity intervention they tend to revert to their less-active lifestyle.¹⁷⁰ Due to the limited number of studies reporting on specific subgroups, we could not perform a comprehensive evaluation of possible sources of heterogeneity, as originally planned.

Our systematic literature review also highlighted a lack of evidence in the literature, particularly in examining the associations of both subjectively and objectively measured sedentary behaviour with outcomes following colorectal cancer, as well as for non-recreational physical activity domains. To enhance our understanding of the impact of both physical activity and sedentary behaviour on colorectal cancer survival, future research should prioritise the assessment of these currently understudied areas.

In consideration of the current evidence base and its methodological limitations, the CUP Global Expert Panel conservatively categorised most of the evidence as limited-no conclusion. Despite the indication of associations across the breadth of the performed analyses, the current evidence is not sufficient to support robust conclusions on causality. The inverse association between recreational activity and all-cause mortality was concluded as limited-suggestive despite being subject to the same biases as the remaining associations, due to the high number of studies providing consistent results (10/11 supported an inverse association, 1 indicative), which was further supported by the results of the linear dose-response meta-analysis. In most cases dose-response meta-analyses could not be performed due to low number of studies, lack of information, and use of a mixture of domains and dimensions assessed, only high versus low levels of activity were compared utilising heterogeneous cut-offs

across studies. Previous assessments, such as those from the ACSM²² that considered the effect of activity across different types of cancer survivors, have concluded a protective effect of physical activity on cancer mortality, albeit with unknown dose required for tangible benefits. However, the CUP Global Expert Panel, focusing specifically on the benefit of activity on survival outcomes of colorectal cancer patients, acknowledged the limitations of the current literature on the field and concluded that the evidence lacks a consistent basis for further recommendations for this specific population. An overview of limitations of cancer survival studies and future research recommendations is presented in the summary manuscript, Box 1³⁰ of the current manuscript series on colorectal cancer survivors.

Cancer survivors as well as their social network members are open to receiving lifestyle advice from healthcare providers.¹⁷¹⁻¹⁷³ The CUP Global Expert Panel recognises that the limitations of the evidence represent an opportunity for further research to clarify the nature of the associations between measures of physical activity and sedentary behaviour and cancer-related outcomes. Furthermore, the panel recognises a need to better inform cancer patients about the links between physical activity and sedentariness and cancer survivorship, beyond making firm recommendations based on high quality evidence.

5 | CONCLUSIONS

Our work indicates that the accumulated evidence on the putative beneficial association of physical activity with outcomes after colorectal cancer is still limited and impacted by methodological limitations frequently present in observational survivorship studies, such as measurement error, residual confounding, reverse causation, and survival bias. As physical activity is a multi-faceted exposure, additional observational research using standardised assessment and ideally repeated objective measures, is of utmost importance to minimise biases and accurately capture its relationship with outcomes after colorectal cancer. Moreover, future high-quality trials focusing on specific physical activity dimensions and domains are essential for formulating robust conclusions, which could form the base for specific lifestyle guidance for colorectal cancer survivors. Nevertheless, even in the absence of strong evidence, there is potential to use this evidence as guidance for colorectal cancer patients and their health professionals to show that a physically active lifestyle and avoidance of sedentary behaviour may be associated with longer overall survival after a colorectal cancer diagnosis.

AUTHOR CONTRIBUTIONS

Konstantinos K. Tsilidis and Doris S. M. Chan are co-principal investigators of CUP Global at Imperial College London (ICL). Doris S. M. Chan and Konstantinos K. Tsilidis implemented the study according to the protocol reviewed by the CUP Global Protocol Expert Group (PEG). Katia Balducci and Sonia Kiss performed the literature search. Katia Balducci, Nerea Becerra-Tomás, Margarita Cariolou, Sonia Kiss, and Rita Vieira performed the study selection and data extraction. Georgios Markozannes performed the study selection, data extraction, and checked,

analysed, and interpreted the data. Dagfinn Aune was a WCRF International CUP Global ICL team member who revised the manuscript. Darren C. Greenwood was a statistical adviser. Amanda J Cross was a CUP Global advisor at Imperial College London. Marc J. Gunter was a CUP Global collaborator on biological processes and provided input into the biological mechanism citations in the manuscript. Ellen Copson was a PEG member, Chair of CUP Global Expert Committee on Cancer Survivorship, and Expert Panel member. Wendy Demark-Wahnefried and Galina Velikova were PEG, OACD, and CUP Global Expert Committee members. Andrew G. Renehan was a PEG member and Deputy Chair of CUP Global Expert Committee on Cancer Survivorship. John Krebs, Matty P. Weijenberg, Monica L. Baskin, Sarah J. Lewis, Jaap Seidell, Rajiv Chowdhury, and Lynette Hill were CUP Global Expert Panel members. Anne M. May, Anne Tjonneland, Karen Steindorf, Martijn Bours, Melissa M. Hudson, Roderick Skinner, and Folakemi T. Odedina were CUP Global Expert Committee members. All members of the CUP Global Expert Committee and Expert Panel provided input into the judgements on the evidence and advised on the interpretation of the review, the public representative (Lynette Hill) did not contribute to the final decisions made by the Panel. Georgios Markozannes drafted the original manuscript. All authors reviewed and provided comments on the manuscript. Doris S. M. Chan is the guarantor and has full access to all the data and takes responsibility for the integrity of the data and the accuracy of the data analysis. The work reported in the paper has been performed by the authors, unless clearly specified in the text.

AFFILIATIONS

¹Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, UK

²Department of Hygiene and Epidemiology, University of Ioannina Medical School, Ioannina, Greece

³Department of Nutrition, Oslo New University College, Oslo, Norway

⁴Department of Research, The Cancer Registry of Norway, Oslo, Norway

⁵Leeds Institute for Data Analytics, Faculty of Medicine and Health, University of Leeds, Leeds, UK

⁶Nutrition and Metabolism Branch, International Agency for Research on Cancer, World Health Organization, Lyon, France

⁷Cancer Sciences Academic Unit, Faculty of Medicine, University of Southampton, Southampton, UK

⁸The Christie NHS Foundation Trust, Manchester Cancer Research Centre, NIHR Manchester Biomedical Research Centre, Division of Cancer Sciences, School of Medical Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester, UK

⁹Department of Epidemiology, GROW School for Oncology and Reproduction, Maastricht University, Maastricht, The Netherlands

¹⁰O'Neal Comprehensive Cancer Center, University of Alabama at Birmingham, Birmingham, Alabama, USA

¹¹Department of Oncology, St Jude Children's Research Hospital, Memphis, Tennessee, USA

¹²Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands

¹³Mayo Clinic Comprehensive Cancer Center, Jacksonville, Florida, USA

¹⁴Department of Paediatric and Adolescent Haematology/Oncology, Great North Children's Hospital and Translational and Clinical Research Institute, and Centre for Cancer, Newcastle University, Newcastle upon Tyne, UK

¹⁵Division of Physical Activity, Prevention and Cancer, German Cancer Research Center (DKFZ) and National Center for Tumor Diseases (NCT), Heidelberg, Germany

¹⁶Danish Cancer Society Research Center, Diet, Cancer and Health, Copenhagen, Denmark

¹⁷Department of Public Health, University of Copenhagen, Copenhagen, Denmark

¹⁸School of Medicine, Faculty of Medicine and Health, University of Leeds, Leeds, UK

¹⁹UPMC Hillman Cancer Center, Pittsburgh, Pennsylvania, USA

²⁰Department of Global Health, Robert Stempel College of Public Health and Social Work, Florida International University, Miami, Florida, USA

²¹World Cancer Research Fund International, London, UK

²²Department of Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

²³Department of Health Sciences, Faculty of Science, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

²⁴Department of Biology, University of Oxford, Oxford, UK

ACKNOWLEDGEMENTS

We thank Teresa Norat for leading the WCRF/AICR Continuous Update Project (CUP) as principal investigator from 2007 to 2020 and for developing the original review protocol. We thank the WCRF International CUP Global ICL team members: Leila Abar, Neesha Nanu, Jakub Sobiecki, Britta Talumaa, and Victoria White, for their contribution to the literature search, study selection, and/or data extraction; and database managers: Rui Vieira, Christophe Stevens, Yusuf O. Anifowoshe, and Lam Teng for implementing and updating the CUP Global database. We also acknowledge the input from the CUP Global Secretariat members: Helen Croker, Panagiota Mitrou, Martin Wiseman, Kate Allen, Nigel T Brockton, Vanessa Gordon-Dseagu (and previous CUP Global Secretariat member Nicole Musuwo), for providing overall coordination for the work and for convening and facilitating discussions with the CUP Global Expert Committee and Expert Panel. We acknowledge the contribution of the Protocol Expert Group (PEG) (Annie Anderson, Steven Clinton, John Mathers, Anne McTiernan, Lesley Turner, and Fränzel van Duijnhooven) and the Outcomes After a Cancer Diagnosis (OACD) CUP Transition workstream (Anne McTiernan, Steven Clinton, Viv Lund) into the development of the original protocol for this work.

FUNDING INFORMATION

This work was funded by the World Cancer Research Fund network of charities (American Institute for Cancer Research [AICR]; World Cancer Research Fund [WCRF]; Wereld Kanker Onderzoek Fonds [WKOF]) (CUP Global Special Grant 2018). The funders of this study

had no role in the decisions about the design and conduct of the study; collection, management, analysis, or interpretation of the data; or the preparation, review, or approval of the manuscript. The process used was based on the method developed by WCRF International's Methodology Task Force for the WCRF/AICR Second Expert Report. The views expressed in this review are the opinions of the authors. They may differ from those in future updates of the evidence related to food, nutrition, physical activity, and cancer incidence and survival.

CONFLICT OF INTEREST STATEMENT

Ellen Copson declared research support from Seca. Galina Velikova declared honoraria from Pfizer, Novartis, and Eisai, an institutional grant from Pfizer, and advisory board and consultancy fees from AstraZeneca, Roche, Novartis, Pfizer, Seagene, Eisai, and Sanofi. All other authors have no conflict of interest related to this work.

DATA AVAILABILITY STATEMENT

Only publicly available data were used in our study. Data sources and handling of these data are described in the materials and methods section. Further details are available from the corresponding author upon request.

ORCID

Georgios Markozannes  <https://orcid.org/0000-0001-8481-579X>

Dagfinn Aune  <https://orcid.org/0000-0002-4533-1722>

Wendy Demark-Wahnefried  <https://orcid.org/0000-0001-5241-932X>

Konstantinos K. Tsilidis  <https://orcid.org/0000-0002-8452-8472>

Doris S. M. Chan  <https://orcid.org/0000-0002-0198-1897>

REFERENCES

1. Morgan E, Arnold M, Gini A, et al. Global burden of colorectal cancer in 2020 and 2040: incidence and mortality estimates from GLOBOCAN. *Gut*. 2023;72:338-344.
2. Gini A, Jansen EEL, Zielonke N, et al. Impact of colorectal cancer screening on cancer-specific mortality in Europe: a systematic review. *Eur J Cancer*. 2020;127:224-235.
3. Lin JS, Piper MA, Perdue LA, et al. Screening for colorectal cancer: updated evidence report and systematic review for the US preventive services task force. *Jama*. 2016;315:2576-2594.
4. Lin JS, Perdue LA, Henrikson NB, Bean SI, Blasi PR. Screening for colorectal cancer: updated evidence report and systematic review for the US preventive services task force. *Jama*. 2021;325:1978-1998.
5. Allemani C, Matsuda T, Di Carlo V, et al. Global surveillance of trends in cancer survival 2000-14 (CONCORD-3): analysis of individual records for 37 513 025 patients diagnosed with one of 18 cancers from 322 population-based registries in 71 countries. *Lancet*. 2018;391:1023-1075.
6. American Cancer Society—Colorectal Cancer Survival Rates. Accessed March 2023. <https://www.cancer.org/cancer/colorectal-cancer/detection-diagnosis-staging/survival-rates.html>
7. Araghi M, Arnold M, Rutherford MJ, et al. Colon and rectal cancer survival in seven high-income countries 2010-2014: variation by age and stage at diagnosis (the ICBP SURVMARK-2 project). *Gut*. 2021;70:114-126.
8. Soerjomataram I, Cbasag C, Bardot A, et al. Cancer survival in Africa, central and south America, and Asia (SURVCAN-3): a

- population-based benchmarking study in 32 countries. *Lancet Oncol.* 2023;24:22-32.
9. World Cancer Research Fund/American Institute for Cancer Research. *Diet, Nutrition, Physical Activity and Cancer: A Global Perspective*. Continuous Update Project Expert Report. 2018. Accessed March 2023. <http://dietandcancerreport.org>
 10. Rezende LFM, Sá TH, Markozannes G, et al. Physical activity and cancer: an umbrella review of the literature including 22 major anatomical sites and 770 000 cancer cases. *Br J Sports Med.* 2018;52:826-833.
 11. McTiernan A, Friedenreich CM, Katzmarzyk PT, et al. Physical activity in cancer prevention and survival: a systematic review. *Med Sci Sports Exerc.* 2019;51:1252-1261.
 12. Je Y, Jeon JY, Giovannucci EL, Meyerhardt JA. Association between physical activity and mortality in colorectal cancer: a meta-analysis of prospective cohort studies. *Int J Cancer.* 2013;133:1905-1913.
 13. Des Guetz G, Uzzan B, Bouillet T, et al. Impact of physical activity on cancer-specific and overall survival of patients with colorectal cancer. *Gastroenterol Res Pract.* 2013;2013:340851.
 14. Schmid D, Leitzmann MF. Association between physical activity and mortality among breast cancer and colorectal cancer survivors: a systematic review and meta-analysis. *Ann Oncol.* 2014;25:1293-1311.
 15. Otto SJ, Korfage IJ, Polinder S, et al. Association of change in physical activity and body weight with quality of life and mortality in colorectal cancer: a systematic review and meta-analysis. *Support Care Cancer.* 2015;23:1237-1250.
 16. Friedenreich CM, Neilson HK, Farris MS, Courneya KS. Physical activity and cancer outcomes: a precision medicine approach. *Clin Cancer Res.* 2016;22:4766-4775.
 17. Wu W, Guo F, Ye J, et al. Pre- and post-diagnosis physical activity is associated with survival benefits of colorectal cancer patients: a systematic review and meta-analysis. *Oncotarget.* 2016;7:52095-52103.
 18. Qiu S, Jiang C, Zhou L. Physical activity and mortality in patients with colorectal cancer: a meta-analysis of prospective cohort studies. *Eur J Cancer Prev.* 2020;29:15-26.
 19. Courneya KS, Vardy JL, O'Callaghan CJ, et al. Effects of a structured exercise program on physical activity and fitness in colon cancer survivors: one year feasibility results from the CHALLENGE trial. *Cancer Epidemiol Biomark Prev.* 2016;25:969-977.
 20. Tremblay MS, Aubert S, Barnes JD, et al. Sedentary Behavior Research Network (SBRN)—terminology consensus project process and outcome. *Int J Behav Nutr Phys Act.* 2017;14:75.
 21. Swain CTV, Nguyen NH, Eagles T, et al. Postdiagnosis sedentary behavior and health outcomes in cancer survivors: a systematic review and meta-analysis. *Cancer.* 2020;126:861-869.
 22. Patel AV, Friedenreich CM, Moore SC, et al. American College of Sports Medicine roundtable report on physical activity, sedentary behavior, and cancer prevention and control. *Med Sci Sports Exerc.* 2019;51:2391-2402.
 23. Campbell KL, Winters-Stone KM, Wiskemann J, et al. Exercise guidelines for cancer survivors: consensus statement from international multidisciplinary roundtable. *Med Sci Sports Exerc.* 2019;51:2375-2390.
 24. Rock CL, Thomson CA, Sullivan KR, et al. American Cancer Society nutrition and physical activity guideline for cancer survivors. *CA Cancer J Clin.* 2022;72:230-262.
 25. WHO guidelines on physical activity and sedentary behaviour. Accessed March 2023. <https://www.who.int/publications/i/item/9789240015128>
 26. Physical Activity Guidelines for Americans. 2018. Accessed March 2023. https://health.gov/sites/default/files/2019-09/Physical_Activity_Guidelines_2nd_edition.pdf,
 27. Global Cancer Update Programme (CUP Global). Accessed March 2023. <https://www.wcrf.org/diet-activity-and-cancer/global-cancer-update-programme/about-the-global-cancer-update-programme/>
 28. Becerra-Tomás N, Markozannes G, Cariolou M, et al. Post-diagnosis adiposity and colorectal cancer prognosis: a Global Cancer Update Programme (CUP Global) systematic literature review and meta-analysis. *Int J Cancer.* 2024;155(3):400-425.
 29. Chan DSM, Cariolou M, Markozannes G, et al. Post-diagnosis dietary factors, supplement use and colorectal cancer prognosis: a Global Cancer Update Programme (CUP Global) systematic literature review and meta-analysis. *Int J Cancer.* 2024;155(3):445-470.
 30. Tsilidis KK, Markozannes G, Becerra-Tomás N, et al. Post-diagnosis adiposity, physical activity, sedentary behaviour, dietary factors, supplement use and colorectal cancer prognosis: Global Cancer Update Programme (CUP Global) summary of evidence grading. *Int J Cancer.* 2024;155(3):471-485.
 31. Page MJ, Moher D, Bossuyt PM, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ.* 2021;372:n160.
 32. Chan DSM, Tsilidis K, Becerra-Tomas N, et al. *Global Cancer Update Programme Protocol for the data collection and systematic literature review on the role of diet, body fatness and physical activity on survival after colorectal cancer*. Accessed December 2023. <https://osf.io/r5ud2/registrations>
 33. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials.* 1986;7:177-188.
 34. Orsini N. Weighted mixed-effects dose-response models for tables of correlated contrasts. *Stata J.* 2021;21:320-347.
 35. Royston P. A strategy for modelling the effect of a continuous covariate in medicine and epidemiology. *Stat Med.* 2000;19:1831-1847.
 36. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ.* 1997;315:629-634.
 37. Duval S, Tweedie R. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics.* 2000;56:455-463.
 38. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med.* 2002;21:1539-1558.
 39. Borenstein M, Higgins JP, Hedges LV, Rothstein HR. Basics of meta-analysis: I(2) is not an absolute measure of heterogeneity. *Res Synth Methods.* 2017;8:5-18.
 40. Viechtbauer W, Cheung MW. Outlier and influence diagnostics for meta-analysis. *Res Synth Methods.* 2010;1:112-125.
 41. USDA. *Nutrition Evidence Systematic Review*. Risk of Bias for Nutrition Observational Studies (RoB-NObs) Tool. Accessed November 2022. <https://nesr.usda.gov/sites/default/files/2019-07/RiskOfBiasForNutritionObservationalStudies-RoB-NObs.pdf>
 42. Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ.* 2016;355:i4919.
 43. Adams SV, Ceballos R, Newcomb PA. Quality of life and mortality of long-term colorectal cancer survivors in the Seattle Colorectal Cancer Family Registry. *PLoS One.* 2016;11:e0156534.
 44. Agalar C, Sokmen S, Arslan C, et al. The impact of sarcopenia on morbidity and long-term survival among patients with peritoneal metastases of colorectal origin treated with cytoreductive surgery and hyperthermic intraperitoneal chemotherapy: a 10-year longitudinal analysis of a single-center experience. *Tech Coloproctol.* 2020;24:301-308.
 45. Araújo RF Jr, Lira GA, Guedes HG, et al. Lifestyle and family history influence cancer prognosis in Brazilian individuals. *Pathol Res Pract.* 2013;209:753-757.

46. Arnold M, Charvat H, Freisling H, et al. Adult overweight and survival from breast and colorectal cancer in Swedish women. *Cancer Epidemiol Biomark Prev*. 2019;28:1518-1524.
47. Baker JR, Umesh S, Jenab M, et al. Prediagnostic blood selenium status and mortality among patients with colorectal cancer in Western European populations. *Biomedicine*. 2021;9:9.
48. Campbell PT, Newton CC, Newcomb PA, et al. Association between body mass index and mortality for colorectal cancer survivors: overall and by tumor molecular phenotype. *Cancer Epidemiol Biomark Prev*. 2015;24:1229-1238.
49. Cannioto RA, Dighe S, Mahoney MC, et al. Habitual recreational physical activity is associated with significantly improved survival in cancer patients: evidence from the Roswell Park Data Bank and BioRepository. *Cancer Causes Control*. 2019;30:1-12.
50. Carr PR, Banbury BL, Berndt SI, et al. Association between intake of red and processed meat and survival in patients with colorectal cancer in a pooled analysis. *Clin Gastroenterol Hepatol*. 2019;17:1561-1570.e3.
51. Carr PR, Jansen L, Walter V, et al. Associations of red and processed meat with survival after colorectal cancer and differences according to timing of dietary assessment. *Am J Clin Nutr*. 2016;103:192-200.
52. Cespedes Feliciano EM, Hohensee C, Rosko AE, et al. Association of prediagnostic frailty, change in frailty status, and mortality after cancer diagnosis in the Women's Health Initiative. *JAMA Netw Open*. 2020;3:e2016747.
53. Cespedes Feliciano EM, Prentice RL, Aragaki AK, et al. Methodological considerations for disentangling a risk factor's influence on disease incidence versus postdiagnosis survival: the example of obesity and breast and colorectal cancer mortality in the Women's Health Initiative. *Int J Cancer*. 2017;141:2281-2290.
54. Charvat H, Freisling H, Noh H, et al. Excess body fatness during early to mid-adulthood and survival from colorectal and breast cancer: a pooled analysis of five international cohort studies. *Cancer Epidemiol Biomark Prev*. 2022;31:325-333.
55. Daniel CR, Shu X, Ye Y, et al. Severe obesity prior to diagnosis limits survival in colorectal cancer patients evaluated at a large cancer centre. *Br J Cancer*. 2016;114:103-109.
56. Diao XY, Peng T, Kong FG, et al. Alcohol consumption promotes colorectal cancer by altering intestinal permeability. *Eur Rev Med Pharmacol Sci*. 2020;24:9370-9377.
57. Dik VK, Murphy N, Siersema PD, et al. Prediagnostic intake of dairy products and dietary calcium and colorectal cancer survival—results from the EPIC cohort study. *Cancer Epidemiol Biomark Prev*. 2014;23:1813-1823.
58. Doria-Rose VP, Newcomb PA, Morimoto LM, Hampton JM, Trentham-Dietz A. Body mass index and the risk of death following the diagnosis of colorectal cancer in postmenopausal women (United States). *Cancer Causes Control*. 2006;17:63-70.
59. Fedirko V, Riboli E, Tjønneland A, et al. Prediagnostic 25-hydroxyvitamin D, VDR and CASR polymorphisms, and survival in patients with colorectal cancer in western European populations. *Cancer Epidemiol Biomark Prev*. 2012;21:582-593.
60. Fedirko V, Romieu I, Aleksandrova K, et al. Pre-diagnostic anthropometry and survival after colorectal cancer diagnosis in Western European populations. *Int J Cancer*. 2014;135:1949-1960.
61. Galas A, Kulig J. Low-grade dietary-related inflammation and survival after colorectal cancer surgery. *J Cancer Res Clin Oncol*. 2014;140:1517-1525.
62. Gibbs DC, Bostick RM, McCullough ML, et al. Association of pre-diagnostic vitamin D status with mortality among colorectal cancer patients differs by common, inherited vitamin D-binding protein isoforms. *Int J Cancer*. 2020;147:2725-2734.
63. Gibson TM, Park Y, Robien K, et al. Body mass index and risk of second obesity-associated cancers after colorectal cancer: a pooled analysis of prospective cohort studies. *J Clin Oncol*. 2014;32:4004-4011.
64. Hardikar S, Newcomb PA, Campbell PT, et al. Prediagnostic physical activity and colorectal cancer survival: overall and stratified by tumor characteristics. *Cancer Epidemiol Biomark Prev*. 2015;24:1130-1137.
65. Haydon AM, Macinnis RJ, English DR, Giles GG. Effect of physical activity and body size on survival after diagnosis with colorectal cancer. *Gut*. 2006;55:62-67.
66. Haydon AM, Macinnis RJ, English DR, Morris H, Giles GG. Physical activity, insulin-like growth factor 1, insulin-like growth factor binding protein 3, and survival from colorectal cancer. *Gut*. 2006;55:689-694.
67. Jacobs S, Harmon BE, Ollberding NJ, et al. Among 4 diet quality indexes, only the alternate Mediterranean diet score is associated with better colorectal cancer survival and only in African American women in the multiethnic cohort. *J Nutr*. 2016;146:1746-1755.
68. Jayasekara H, English DR, Haydon A, et al. Associations of alcohol intake, smoking, physical activity and obesity with survival following colorectal cancer diagnosis by stage, anatomic site and tumor molecular subtype. *Int J Cancer*. 2018;142:238-250.
69. Je Y, Lee JE, Ma J, et al. Prediagnostic plasma vitamin B6 (pyridoxal 5'-phosphate) and survival in patients with colorectal cancer. *Cancer Causes Control*. 2013;24:719-729.
70. Jeffreys M, Redaniel MT, Martin RM. The effect of pre-diagnostic vitamin D supplementation on cancer survival in women: a cohort study within the UK Clinical Practice Research Datalink. *BMC Cancer*. 2015;15:670.
71. Kocarnik JM, Chan AT, Slattery ML, et al. Relationship of prediagnostic body mass index with survival after colorectal cancer: stage-specific associations. *Int J Cancer*. 2016;139:1065-1072.
72. Kyrø C, Frederiksen K, Holm M, et al. Prediagnosis plasma concentrations of enterolactone and survival after colorectal cancer: the Danish Diet, Cancer and Health cohort. *Br J Nutr*. 2019;122:552-563.
73. Laake I, Larsen IK, Selmer R, Thune I, Veierød MB. Pre-diagnostic body mass index and weight change in relation to colorectal cancer survival among incident cases from a population-based cohort study. *BMC Cancer*. 2016;16:402.
74. Lewis C, Xun P, Fly AD, Luo J, He K. Fish oil supplementation and quality of life in stage II colorectal cancer patients: a 24-month follow-up study. *Nutr Cancer*. 2015;67:1239-1246.
75. Mao Z, Aglago EK, Zhao Z, et al. Dietary intake of advanced glycation end products (AGEs) and mortality among individuals with colorectal cancer. *Nutrients*. 2021;13:4435.
76. Ng K, Meyerhardt JA, Wu K, et al. Circulating 25-hydroxyvitamin d levels and survival in patients with colorectal cancer. *J Clin Oncol*. 2008;26:2984-2991.
77. Ogino S, Shima K, Nosho K, et al. A cohort study of p27 localization in colon cancer, body mass index, and patient survival. *Cancer Epidemiol Biomark Prev*. 2009;18:1849-1858.
78. Oyeyemi SO, Braaten T, Skeie G, Borch KB. Competing mortality risks analysis of prediagnostic lifestyle and dietary factors in colorectal cancer survival: the Norwegian Women and Cancer Study. *BMJ Open Gastroenterol*. 2019;6:e000338.
79. Park SM, Lim MK, Shin SA, Yun YH. Impact of prediagnosis smoking, alcohol, obesity, and insulin resistance on survival in male cancer patients: National Health Insurance Corporation Study. *J Clin Oncol*. 2006;24:5017-5024.
80. Park SM, Yun YH, Kim YA, et al. Prediagnosis body mass index and risk of secondary primary cancer in male cancer survivors: a large cohort study. *J Clin Oncol*. 2016;34:4116-4124.
81. Patel SA, Chen YH, Hornick JL, et al. Early-stage rectal cancer: clinical and pathologic prognostic markers of time to local recurrence and overall survival after resection. *Dis Colon Rectum*. 2014;57:449-459.
82. Pelsler C, Arem H, Pfeiffer RM, et al. Prediagnostic lifestyle factors and survival after colon and rectal cancer diagnosis in the National Institutes of Health (NIH)-AARP Diet and Health Study. *Cancer*. 2014;120:1540-1547.

83. Prizment AE, Flood A, Anderson KE, Folsom AR. Survival of women with colon cancer in relation to precancer anthropometric characteristics: the Iowa Women's Health Study. *Cancer Epidemiol Biomark Prev.* 2010;19:2229-2237.
84. Reichle K, Peter RS, Concin H, Nagel G. Associations of pre-diagnostic body mass index with overall and cancer-specific mortality in a large Austrian cohort. *Cancer Causes Control.* 2015;26:1643-1652.
85. Romaguera D, Ward H, Wark PA, et al. Pre-diagnostic concordance with the WCRF/AICR guidelines and survival in European colorectal cancer patients: a cohort study. *BMC Med.* 2015;13:107.
86. Sharma I, Roebroth B, Zhu Y, et al. Hypothesis and data-driven dietary patterns and colorectal cancer survival: findings from Newfoundland and Labrador colorectal cancer cohort. *Nutr J.* 2018;17:55.
87. Skeie G, Braaten T, Olsen A, et al. Whole grain intake and survival among Scandinavian colorectal cancer patients. *Nutr Cancer.* 2014;66:6-13.
88. Slattery ML, Anderson K, Samowitz W, et al. Hormone replacement therapy and improved survival among postmenopausal women diagnosed with colon cancer (USA). *Cancer Causes Control.* 1999;10:467-473.
89. Slattery ML, French TK, Egger MJ, Lyon JL. Diet and survival of patients with colon cancer in Utah: is there an association? *Int J Epidemiol.* 1989;18:792-797.
90. Stapley S, Peters TJ, Sharp D, Hamilton W. The mortality of colorectal cancer in relation to the initial symptom at presentation to primary care and to the duration of symptoms: a cohort study using medical records. *Br J Cancer.* 2006;95:1321-1325.
91. Sun H, Liu Y, Huang H, Li D, Zhao Y. Diet quality score and survival rate in patients with colorectal cancer. *Asia Pac J Clin Nutr.* 2019;28:601-606.
92. Walter V, Jansen L, Knebel P, Chang-Claude J, Hoffmeister M, Brenner H. Physical activity and survival of colorectal cancer patients: population-based study from Germany. *Int J Cancer.* 2017;140:1985-1997.
93. Walter V, Jansen L, Ulrich A, et al. Alcohol consumption and survival of colorectal cancer patients: a population-based study from Germany. *Am J Clin Nutr.* 2016;103:1497-1506.
94. Wang N, Khankari NK, Cai H, et al. Prediagnosis body mass index and waist-hip circumference ratio in association with colorectal cancer survival. *Int J Cancer.* 2017;140:292-301.
95. Ward HA, Norat T, Overvad K, et al. Pre-diagnostic meat and fibre intakes in relation to colorectal cancer survival in the European Prospective Investigation into Cancer and Nutrition. *Br J Nutr.* 2016;116:316-325.
96. Weinstein SJ, Mondul AM, Yu K, et al. Circulating 25-hydroxyvitamin D up to 3 decades prior to diagnosis in relation to overall and organ-specific cancer survival. *Eur J Epidemiol.* 2018;33:1087-1099.
97. Wolpin BM, Wei EK, Ng K, et al. Prediagnostic plasma folate and the risk of death in patients with colorectal cancer. *J Clin Oncol.* 2008;26:3222-3228.
98. Xia S, Sun M, Liu X. Major depression but not minor to intermediate depression correlates with unfavorable prognosis in surgical colorectal cancer patients underwent adjuvant chemotherapy. *Psychol Health Med.* 2020;25:309-318.
99. Yang Y, Mauldin PD, Ebeling M, et al. Effect of metabolic syndrome and its components on recurrence and survival in colon cancer patients. *Cancer.* 2013;119:1512-1520.
100. Yuan C, Song M, Zhang Y, et al. Prediagnostic circulating concentrations of vitamin D binding protein and survival among patients with colorectal cancer. *Cancer Epidemiol Biomark Prev.* 2020;29:2323-2331.
101. Zamora-Ros R, Guinó E, Alonso MH, et al. Dietary flavonoids, lignans and colorectal cancer prognosis. *Sci Rep.* 2015;5:14148.
102. Zell JA, Lin BS, Ziogas A, Anton-Culver H. Meat consumption, ornithine decarboxylase gene polymorphism, and outcomes after colorectal cancer diagnosis. *J Carcinog.* 2012;11:17.
103. Zell JA, McEligot AJ, Ziogas A, Holcombe RF, Anton-Culver H. Differential effects of wine consumption on colorectal cancer outcomes based on family history of the disease. *Nutr Cancer.* 2007;59:36-45.
104. Zell JA, Ziogas A, Bernstein L, et al. Meat consumption, nonsteroidal anti-inflammatory drug use, and mortality among colorectal cancer patients in the California Teachers Study. *Cancer Prev Res (Phila).* 2010;3:865-875.
105. Zhou J, Ge X, Fan X, Wang J, Miao L, Hang D. Associations of vitamin D status with colorectal cancer risk and survival. *Int J Cancer.* 2021;149:606-614.
106. Zhu Y, Zhao J, Vallis J, et al. Prediagnostic consumption of vitamin D, calcium and dairy products and colorectal cancer survival: results from the Newfoundland Colorectal Cancer Registry Cohort Study. *Br J Nutr.* 2022;128:290-299.
107. Aarts MJ, Kamphuis CB, Louwman MJ, Coebergh JW, Mackenbach JP, van Lenthe FJ. Educational inequalities in cancer survival: a role for comorbidities and health behaviours? *J Epidemiol Community Health.* 2013;67:365-373.
108. Annie FH, Uejio CK, Bhagat A, Kochar T, Embrey S, Tager A. Survival analysis of cancer patients of differing payer type in South West Virginia, between 2000 and 2013. *Cureus.* 2018;10:e3022.
109. Boyle T, Fritschi L, Platell C, Heyworth J. Lifestyle factors associated with survival after colorectal cancer diagnosis. *Br J Cancer.* 2013;109:814-822.
110. Gannavarapu BS, Lau SKM, Carter K, et al. Prevalence and survival impact of pretreatment cancer-associated weight loss: a tool for guiding early palliative care. *J Oncol Pract.* 2018;14:e238-e250.
111. Minami Y, Kanemura S, Oikawa T, et al. Associations of Japanese food intake with survival of stomach and colorectal cancer: a prospective patient cohort study. *Cancer Sci.* 2020;111:2558-2569.
112. Phipps AI, Baron J, Newcomb PA. Prediagnostic smoking history, alcohol consumption, and colorectal cancer survival: the Seattle Colon Cancer Family Registry. *Cancer.* 2011;117:4948-4957.
113. Phipps AI, Robinson JR, Campbell PT, et al. Prediagnostic alcohol consumption and colorectal cancer survival: the Colon Cancer Family Registry. *Cancer.* 2017;123:1035-1043.
114. Park J, Kim JH, Lee HJ, et al. The effects of physical activity and body fat mass on colorectal polyp recurrence in patients with previous colorectal cancer. *Cancer Prev Res (Phila).* 2017;10:478-484.
115. Jeon J, Sato K, Niedzwiecki D, et al. Impact of physical activity after cancer diagnosis on survival in patients with recurrent colon cancer: Findings from CALGB 89803/Alliance. *Clin Colorectal Cancer.* 2013;12:233-238.
116. Cheng E, Ou FS, Ma C, et al. Diet- and lifestyle-based prediction models to estimate cancer recurrence and death in patients with stage III colon cancer (CALGB 89803/Alliance). *J Clin Oncol.* 2022;40:740-751.
117. van Zutphen M, van Duijnhoven FJB, Wesseling E, et al. Identification of lifestyle behaviors associated with recurrence and survival in colorectal cancer patients using random survival forests. *Cancer.* 2021;13:2442.
118. Morikawa T, Kuchiba A, Yamauchi M, et al. Association of CTNNB1 (beta-catenin) alterations, body mass index, and physical activity with survival in patients with colorectal cancer. *Jama.* 2011;305:1685-1694.
119. Cheng CC, Lai IL, Huang SH, et al. Association of preoperative physical activity with short- and long-term outcomes in patients undergoing palliative resection for metastatic colorectal cancer: an inverse probability of treatment weighting analysis. *Cancer.* 2022;14:489.

120. You JF, Hsu YJ, Chern YJ, et al. Association of a preoperative leisure-time physical activity with short- and long-term outcomes of patients undergoing curative resection for stage I to III colorectal cancer: a propensity score matching analysis. *Dis Colon Rectum*. 2020;63:796-806.
121. Piringer G, Vormittag L, Ohler L, et al. REGO-ACT: assessment of physical activity during treatment with regorafenib for metastatic colorectal cancer. *Wien Klin Wochenschr*. 2020;132:423-430.
122. Lee M, Lee Y, Jang D, Shin A. Physical activity after colorectal cancer diagnosis and mortality in a Nationwide Retrospective Cohort Study. *Cancer*. 2021;13:4804.
123. van Zutphen M, Boshuizen HC, Kenkhuis MF, et al. Lifestyle after colorectal cancer diagnosis in relation to recurrence and all-cause mortality. *Am J Clin Nutr*. 2021;113:1447-1457.
124. Tarasenko YN, Linder DF, Miller EA. Muscle-strengthening and aerobic activities and mortality among 3+ year cancer survivors in the U.S. *Cancer Causes Control*. 2018;29:475-484.
125. Tamakoshi A, Nakamura K, Ukawa S, et al. Characteristics and prognosis of Japanese colorectal cancer patients: the BioBank Japan Project. *J Epidemiol*. 2017;27:S36-S42.
126. Hanyuda A, Kim SA, Martinez-Fernandez A, et al. Survival benefit of exercise differs by tumor IRS1 expression status in colorectal cancer. *Ann Surg Oncol*. 2016;23:908-917.
127. Ratjen I, Schafmayer C, di Giuseppe R, et al. Postdiagnostic physical activity, sleep duration, and TV watching and all-cause mortality among long-term colorectal cancer survivors: a prospective cohort study. *BMC Cancer*. 2017;17:701.
128. Arem H, Pfeiffer RM, Engels EA, et al. Pre- and postdiagnosis physical activity, television viewing, and mortality among patients with colorectal cancer in the National Institutes of Health-AARP Diet and Health Study. *J Clin Oncol*. 2015;33:180-188.
129. Cao Y, Meyerhardt JA, Chan AT, Wu K, Fuchs CS, Giovannucci EL. Television watching and colorectal cancer survival in men. *Cancer Causes Control*. 2015;26:1467-1476.
130. Campbell PT, Patel AV, Newton CC, Jacobs EJ, Gapstur SM. Associations of recreational physical activity and leisure time spent sitting with colorectal cancer survival. *J Clin Oncol*. 2013;31:876-885.
131. Yamauchi M, Lochhead P, Imamura Y, et al. Physical activity, tumor PTGS2 expression, and survival in patients with colorectal cancer. *Cancer Epidemiol Biomark Prev*. 2013;22:1142-1152.
132. Kuiper JG, Phipps AI, Neuhauser ML, et al. Recreational physical activity, body mass index, and survival in women with colorectal cancer. *Cancer Causes Control*. 2012;23:1939-1948.
133. Baade PD, Meng X, Youl PH, Aitken JF, Dunn J, Chambers SK. The impact of body mass index and physical activity on mortality among patients with colorectal cancer in Queensland, Australia. *Cancer Epidemiol Biomark Prev*. 2011;20:1410-1420.
134. Meyerhardt JA, Ogino S, Kirkner GJ, et al. Interaction of molecular markers and physical activity on mortality in patients with colon cancer. *Clin Cancer Res*. 2009;15:5931-5936.
135. Meyerhardt JA, Giovannucci EL, Ogino S, et al. Physical activity and male colorectal cancer survival. *Arch Intern Med*. 2009;169:2102-2108.
136. Meyerhardt JA, Giovannucci EL, Holmes MD, et al. Physical activity and survival after colorectal cancer diagnosis. *J Clin Oncol*. 2006;24:3527-3534.
137. Smit KC, Derksen JWG, Beets GLO, et al. Physical activity is associated with improved overall survival among patients with metastatic colorectal cancer. *Cancer*. 2022;14:1001.
138. Guercio BJ, Zhang S, Ou FS, et al. Associations of physical activity with survival and progression in metastatic colorectal cancer: results from Cancer and Leukemia Group B (Alliance)/SWOG 80405. *J Clin Oncol*. 2019;37:2620-2631.
139. Van Blarigan EL, Fuchs CS, Niedzwiecki D, et al. Association of survival with adherence to the American Cancer Society nutrition and physical activity guidelines for cancer survivors after colon cancer diagnosis: the CALGB 89803/Alliance trial. *JAMA Oncol*. 2018;4:783-790.
140. Meyerhardt JA, Heseltine D, Niedzwiecki D, et al. Impact of physical activity on cancer recurrence and survival in patients with stage III colon cancer: findings from CALGB 89803. *J Clin Oncol*. 2006;24:3535-3541.
141. Phipps AI, Shi Q, Zemla TJ, et al. Physical activity and outcomes in patients with stage III colon cancer: a correlative analysis of phase III trial NCCTG N0147 (Alliance). *Cancer Epidemiol Biomark Prev*. 2018;27:696-703.
142. Ainsworth BE, Haskell WL, Whitt MC, et al. Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc*. 2000;32:S498-S504.
143. Quadrilatero J, Hoffman-Goetz L. Physical activity and colon cancer. A systematic review of potential mechanisms. *J Sports Med Phys Fitness*. 2003;43:121-138.
144. Wang T, Zhang Y, Taaffe DR, et al. Protective effects of physical activity in colon cancer and underlying mechanisms: a review of epidemiological and biological evidence. *Crit Rev Oncol Hematol*. 2022;170:103578.
145. Sax AT, Jenkins DG, Devin JL, Hughes GI, Bolam KA, Skinner TL. The insulin-like growth factor axis: a biological mechanism linking physical activity to colorectal cancer survival. *Cancer Epidemiol*. 2014;38:455-459.
146. Friedenreich CM, Shaw E, Neilson HK, Brenner DR. Epidemiology and biology of physical activity and cancer recurrence. *J Mol Med (Berl)*. 2017;95:1029-1041.
147. Taylor CA, Cheng CP, Espinosa LA, Tang BT, Parker D, Herfkens RJ. In vivo quantification of blood flow and wall shear stress in the human abdominal aorta during lower limb exercise. *Ann Biomed Eng*. 2002;30:402-408.
148. Regmi S, Fu A, Luo KQ. High shear stresses under exercise condition destroy circulating tumor cells in a microfluidic system. *Sci Rep*. 2017;7:39975.
149. Brown JC, Rhim AD, Manning SL, et al. Effects of exercise on circulating tumor cells among patients with resected stage I-III colon cancer. *PLoS One*. 2018;13:e0204875.
150. Liberman K, Forti LN, Beyer I, Bautmans I. The effects of exercise on muscle strength, body composition, physical functioning and the inflammatory profile of older adults: a systematic review. *Curr Opin Clin Nutr Metab Care*. 2017;20:30-53.
151. Petersen AM, Pedersen BK. The anti-inflammatory effect of exercise. *J Appl Physiol (Bethesda, MD 1985)*. 2005;98:1154-1162.
152. Pedersen BK, Febbraio MA. Muscles, exercise and obesity: skeletal muscle as a secretory organ. *Nat Rev Endocrinol*. 2012;8:457-465.
153. Shim YJ, Kim HJ, Oh SC, Lee SI, Choi SW. Exercise during adjuvant treatment for colorectal cancer: treatment completion, treatment-related toxicities, body composition, and serum level of adipokines. *Cancer Manag Res*. 2019;11:5403-5412.
154. Darkwah S, Park EJ, Myint PK, et al. Potential roles of muscle-derived extracellular vesicles in remodeling cellular microenvironment: proposed implications of the exercise-induced myokine, irisin. *Front Cell Dev Biol*. 2021;9:634853.
155. Zhang Y, Kim JS, Wang TZ, et al. Potential role of exercise induced extracellular vesicles in prostate cancer suppression. *Front Oncol*. 2021;11:746040.
156. Pedersen BK. Exercise-induced myokines and their role in chronic diseases. *Brain Behav Immun*. 2011;25:811-816.
157. McClellan JL, Steiner JL, Day SD, et al. Exercise effects on polyp burden and immune markers in the ApcMin/+ mouse model of intestinal tumorigenesis. *Int J Oncol*. 2014;45:861-868.
158. Barozzi C, Ravaioli M, D'Errico A, et al. Relevance of biologic markers in colorectal carcinoma: a comparative study of a broad panel. *Cancer*. 2002;94:647-657.

159. Glass AR, Kikendall JW, Sobin LH, Bowen PE. Serum concentrations of insulin-like growth factor 1 in colonic neoplasia. *Acta Oncol (Stockh, Swed)*. 1994;33:70-71.
160. Renehan AG, Painter JE, Atkin WS, Potten CS, Shalet SM, O'Dwyer ST. High-risk colorectal adenomas and serum insulin-like growth factors. *Br J Surg*. 2001;88:107-113.
161. Health Education Materials With/Out a Physical Activity Program for Patients Who Have Undergone Treatment for High-Risk Stage II or Stage III Colon Cancer (CHALLENGE). Accessed November 2023. <https://clinicaltrials.gov/study/NCT00819208>
162. Gorzelitz J, Peppard PE, Malecki K, Gennuso K, Nieto FJ, Cadmus-Bertram L. Predictors of discordance in self-report versus device-measured physical activity measurement. *Ann Epidemiol*. 2018;28:427-431.
163. Boyle T, Lynch BM, Courneya KS, Vallance JK. Agreement between accelerometer-assessed and self-reported physical activity and sedentary time in colon cancer survivors. *Support Care Cancer*. 2015;23:1121-1126.
164. Courneya KS, Friedenreich CM. Relationship between exercise pattern across the cancer experience and current quality of life in colorectal cancer survivors. *J Altern Complement Med (New York, NY)*. 1997;3:215-226.
165. Cabilan CJ, Hines S. The short-term impact of colorectal cancer treatment on physical activity, functional status and quality of life: a systematic review. *JBI Database System Rev Implement Rep*. 2017;15:517-566.
166. Adamsen L, Andersen C, Lill Lund C, Bloomquist K, Møller T. Rethinking exercise identity: a qualitative study of physically inactive cancer patients' transforming process while undergoing chemotherapy. *BMJ Open*. 2017;7:e016689.
167. Renman D, Strigård K, Palmqvist R, Näsvalld P, Gunnarsson U, Edin-Liljgren A. Attitudes to and experiences of physical activity after colon cancer diagnosis amongst physically active individuals—a qualitative study. *Cancer Control*. 2022;29:10732748221119352.
168. Mosher CE, Sloane R, Morey MC, et al. Associations between life-style factors and quality of life among older long-term breast, prostate, and colorectal cancer survivors. *Cancer*. 2009;115:4001-4009.
169. Mols F, Beijers AJ, Vreugdenhil G, Verhulst A, Schep G, Husson O. Chemotherapy-induced peripheral neuropathy, physical activity and health-related quality of life among colorectal cancer survivors from the PROFILES registry. *J Cancer Surviv Res Pract*. 2015;9:512-522.
170. Husson O, Mols F, Ezendam NP, Schep G, van de Poll-Franse LV. Health-related quality of life is associated with physical activity levels among colorectal cancer survivors: a longitudinal, 3-year study of the PROFILES registry. *J Cancer Surviv Res Pract*. 2015;9:472-480.
171. Williams K, Beeken RJ, Wardle J. Health behaviour advice to cancer patients: the perspective of social network members. *Br J Cancer*. 2013;108:831-835.
172. Beeken RJ, Williams K, Wardle J, Croker H. "What about diet?" A qualitative study of cancer survivors' views on diet and cancer and their sources of information. *Eur J Cancer Care*. 2016;25:774-783.
173. Smith L, Croker H, Fisher A, Williams K, Wardle J, Beeken RJ. Cancer survivors' attitudes towards and knowledge of physical activity, sources of information, and barriers and facilitators of engagement: a qualitative study. *Eur J Cancer Care*. 2017;26:e12641.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Markozannes G, Becerra-Tomás N, Cariolou M, et al. Post-diagnosis physical activity and sedentary behaviour and colorectal cancer prognosis: A Global Cancer Update Programme (CUP Global) systematic literature review and meta-analysis. *Int J Cancer*. 2024;155(3):426-444. doi:10.1002/ijc.34903