

Brief Report

Association Between the PRIMARY Score at Staging Prostate-specific Membrane Antigen Positron Emission Tomography and Overall Survival Among Patients with Newly Diagnosed Prostate Cancer: Findings from the International, Multicenter PROMISE Registry

Madeleine J. Karpinski^{a,b,*}, Sebastian Hoferück^c, Wolfgang P. Fendler^a, Caner Civan^a, Ralph A. Bundschuh^c, Christian Thomas^d, Anders Bjartell^{e,f}, Elin Trägårdh^{f,g,h}, Timo F.W. Soeterik^{ij}, Laura Evangelista^{k,l}, Andrej Vondrak^m, Sazan Rasulⁿ, Laura Forner^a, Andrea Di Giorgio^o, Helen Scholtissek^p, Jonathan Miksch^{q,r}, Adrien Holzgreve^{s,t}, Francesco Lanfranchi^u, Kambiz Rahbar^b, Michael S. Hofman^{v,w}, Isabel Rauscher^x, Osman Güven^x, Matthias Eiber^x, Narjess Ayati^y, Lale Umutlu^z, Ken Herrmann^a, Boris Hadaschik^{aa,†}, Louise Emmett^{y,†}

^a Department of Nuclear Medicine, DKTK and NCT University Hospital Essen, Essen, Germany; ^b Department of Nuclear Medicine, University Hospital Münster, Münster, Germany; ^c Department of Nuclear Medicine, University Hospital Carl Gustav Carus, TU Dresden, Dresden, Germany; ^d Department of Urology, University Hospital Carl Gustav Carus, TU Dresden, Dresden, Germany; ^e Department of Urology, Skåne University Hospital, Malmö, Sweden; ^f Department of Translational Medicine, Lund University, Malmö, Sweden; ^g Department of Clinical Physiology and Nuclear Medicine, Skåne University Hospital, Malmö, Sweden; ^h Wallenberg Centre of Molecular Medicine, Lund University, Lund, Sweden; ⁱ Department of Radiation Oncology, University Medical Center Utrecht, Utrecht, The Netherlands; ^j Department of Urology, St. Antonius Hospital, Nieuwegein/Utrecht, The Netherlands; ^k Department of Biomedical Sciences, Humanitas University, Pieve Emanuele, Italy; ^l Nuclear Medicine Unit, IRCCS Humanitas Research Hospital, Rozzano, Italy; ^m Nuclear Medicine Department, IZOTOPCENTRUM s.r.o, Nitra, Slovakia; ⁿ Department of Biomedical Imaging and Image-guided Therapy, Medical University of Vienna, Vienna, Austria; ^o Department of Nuclear Medicine, Alma Mater Studiorum, University of Bologna, Bologna, Italy; ^p Department of Nuclear Medicine, University of Augsburg, Augsburg, Germany; ^q Department of Nuclear Medicine, Ulm University Hospital, Ulm, Germany; ^r Department of Radiology, Ulm University Hospital, Ulm, Germany; ^s Department of Nuclear Medicine, LMU University Hospital, LMU Munich, Munich, Germany; ^t Ahmanson Translational Theranostics Division, UCLA David Geffen School of Medicine, Los Angeles, CA, USA; ^u Department of Experimental Medicine, University of Genoa, Genoa, Italy; ^v Prostate Cancer Theranostics and Imaging Centre of Excellence, Peter MacCallum Centre Melbourne, Australia; ^w Sir Peter MacCallum Department of Oncology, University of Melbourne, Melbourne, Australia; ^x Department of Nuclear Medicine, School of Medicine and Health, TUM University Hospital, Munich, Germany; ^y Department of Theranostics and Nuclear Medicine, St. Vincent's Hospital, Sydney, Australia; ^z Department of Diagnostic and Interventional Radiology and Neuroradiology, University Hospital Essen, Essen, Germany; ^{aa} Department of Urology, DKTK and NCT University Hospital Essen, Essen, Germany

Article info

Article history:

Accepted October 13, 2025

Associate Editor:

Gianluca Giannarini, MD

Abstract

The PRIMARY score was implemented in Prostate Cancer Molecular Imaging Standardized Evaluation (PROMISE) version 2 to improve accuracy for the diagnosis of clinically significant prostate cancer using prostate-specific membrane antigen (PSMA) positron emission tomography (PET). We reviewed overall survival (OS) for patients who underwent PSMA PET for initial staging to evaluate the prognostic value of PRIMARY in a large, international, multicenter cohort. The cohort comprised 1889

[†] These authors contributed equally and share senior authorship.

* Corresponding author. Department of Nuclear Medicine, University Hospital Essen, Hufelandstrasse 55, 45147 Essen, Germany. Tel. +49 201 723 2032.

E-mail address: madeleine.karpinski@uk-essen.de (M.J. Karpinski).

Keywords:

Overall survival;
PRIMARY score;
Prostate cancer;
Prostate-specific membrane
antigen;
Positron emission tomography

patients who underwent PSMA PET for initial staging of prostate cancer at investigator sites across Europe and Australia between 2012 and 2021. Hazard ratios (HRs) with 95% confidence interval (CI) were calculated for PRIMARY scores to identify predictors of OS. Complete-case head-to-head comparisons were conducted for Prostate Imaging-Reporting and Data System (PI-RADS) versus PRIMARY scores, and cT stage versus PRIMARY scores. We present preliminary findings up to January 31, 2025, when 231 deaths had occurred. PRIMARY score 5 (HR 1.5, 95% CI 1.0–2.3; $p = 0.045$) was associated with shorter OS. Improvements in C index values confirmed the added prognostic value of the PRIMARY score when combined with PI-RADS or cT stage. PRIMARY score 5 on initial PSMA PET is prognostic for shorter OS. There is ongoing long-term follow-up in the PROMISE registry (NCT06320223, promise-pet.org).

© 2025 The Author(s). Published by Elsevier B.V. on behalf of European Association of Urology. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

ADVANCING PRACTICE**What does this study add?**

For the first time we confirm the prognostic value of PROMISE and PRIMARY using a large international multicenter cohort of patients who underwent prostate-specific membrane antigen (PSMA) positron emission tomography (PET) at initial staging.

Clinical Relevance

This large international registry analysis provides the first evidence that a PRIMARY score of 5 on initial staging PSMA PET is independently associated with shorter overall survival in patients with prostate cancer, offering additive prognostic value when combined with PI-RADS or clinical T stage. However, the results should be interpreted with caution given the retrospective design, heterogeneous imaging protocols, and relatively short follow-up, which may limit generalizability. Nevertheless, these findings support the potential role of standardized PSMA-PET metrics in refining risk stratification and informing treatment strategies in patients with prostate cancer at the time of first diagnosis. Associate Editor: Gianluca Giannarini, MD.

Patient Summary

We looked at the value of tools called the PROMISE criteria and the PRIMARY score in predicting the prognosis for patients with prostate cancer who undergo a newer type of scan called PSMA PET (prostate-specific membrane antigen positron emission tomography).

The standard for diagnosis of prostate cancer (PC) in men is multiparametric magnetic resonance imaging (mpMRI) in accordance with the Prostate Imaging-Reporting and Data System (PI-RADS), and subsequent biopsy [1]. The additive diagnostic accuracy of prostate-specific membrane antigen (PSMA) positron emission tomography (PET) to mpMRI has been evaluated [2,3]. On the basis of its practical value, the 5-point PRIMARY score, developed to describe local PSMA PET findings and optimize cancer diagnosis [4], was implemented in Prostate Cancer Molecular Imaging Standardized Evaluation (PROMISE) v2 [5]. The PRIMARY score combines the intraprostatic pattern and site and PSMA activity. Score 1 (no intraprostatic pattern) and score 2 (diffuse transition-zone or symmetric central-zone activity without focal uptake) are deemed “benign” patterns, while score 3 (focal transition-zone activity two times higher than the background), score 4 (focal peripheral-zone activity), and score 5 (maximum standardized uptake value [SUV_{max}]

≥ 12) are deemed “malignant” patterns [4]. When combined with mpMRI, PSMA PET significantly improves the sensitivity and negative predictive value for detection of clinically significant PC (csPC) over mpMRI alone [6]. However, to the best of our knowledge, the PRIMARY score has not yet been associated with survival in a large, international, multicenter cohort. Here we report on the prognostic value of the PRIMARY score for overall survival (OS) in a primary staging subgroup from the PROMISE registry study (ClinicalTrials.gov NCT06320223).

We present preliminary findings for a subgroup of patients at initial staging ($n = 1889$). Inclusion criteria and central data collection for the PROMISE registry have previously been described [7] and are presented in more detail in the [Supplementary material](#).

Patient data, including laboratory parameters, histopathology, clinical data, and conventional imaging reports according to local site practice, were screened to

classify patients in an initial staging subgroup. The patients included had not received definitive primary therapy before PSMA PET imaging.

All patients underwent ⁶⁸Ga-based or ¹⁸F-based PSMA PET-computed tomography (CT) or whole-body PET-MRI between December 6, 2012 and October 15, 2021 for initial staging of PC.

In alignment with the PROMISE v2 criteria [5], PRIMARY scores were retrospectively assigned on the basis of PSMA PET images and clinical PET reports. Data for clinical parameters (Gleason grade group, cT stage, and prostate-specific antigen [PSA] at the time of PSMA PET) were collected when available. PI-RADS scores for MRI results were collected when available.

The primary endpoint of OS was defined as time from PSMA PET to death from any cause. Follow-up was calculated as the time from PSMA PET to the last day the patient was known to be alive. Vital status for patients was requested from ongoing therapy schedules, electronic medical records, or cancer registry data sets according to local site practice.

Cox proportional-hazard ratios (HRs) with 95% confidence interval (CI) were calculated for the PRIMARY score. Multivariable Cox regression was performed for patients with complete data sets for Gleason grade group, PSA level at PET, and cT stage at diagnosis (complete-case analysis).

Head-to-head subgroup analyses were performed for patients with complete data for PI-RADS and PRIMARY scores (*n* = 366), and for patients with cT stage and PRI-

MARY scores (*n* = 960). cT stage was defined on the basis of clinical results without PET. Separate Cox proportional-hazard models were built for the PRIMARY score, PI-RADS score, and cT stage. The additive value of the PRIMARY score when combined with either the PI-RADS score or cT stage was measured in terms of the C index for combined Cox proportional-hazard models. Internal bootstrap validation was conducted by running 1000 bootstrap replicates for the combined models. We used R v4.4.1 (R Foundation for Statistical Computing, Vienna, Austria) for all statistical analyses, with statistical significance set at *p* < 0.05.

Baseline characteristics of the initial staging subgroup are presented in [Supplementary Table 1](#). In total, 231 patients had died by January 31, 2025. The majority of patients had PRIMARY score 4 (*n* = 642, 34%) or score 5 (*n* = 810, 43%); 437 patients (23%) had PRIMARY score 1–3.

Univariable regression analysis revealed that PRIMARY score 5, miT3b/T4 stage, and all PROMISE categories of locoregional or distant disease were predictors for shorter OS ([Fig. 1](#)).

Multivariable regression results (*n* = 925) are presented in [Supplementary Fig. 1](#). PRIMARY score 3 and stages cT2 and cT4 were predictors for worse OS, while Gleason grade group and PSA were risk factors, but did not reach statistical significance.

We tested the additive value of the PRIMARY score when combined with the PI-RADS score or with cT stage ([Table 1](#)). The C index values were 0.68 (95% CI 0.58–0.77) for PI-RADS + PRIMARY, and 0.65 (95% CI 0.59–0.71) for cT stage + PRI-




Fig. 1 – Univariable regression results for PROMISE and PSMA PET metrics in the initial staging subgroup (*n* = 1889) for identification of potential predictors of overall survival. Estimates are the hazard ratio with 95% confidence interval. T, N, and M stages refer to the molecular imaging TNM classification (miTNM). PROMISE = Prostate Cancer Molecular Imaging Standardized Evaluation; PSMA = prostate-specific membrane antigen; PET = positron emission tomography.

Table 1 – Additive predictive value of the PRIMARY score in separate combined models with PI-RADS or cT stage (complete-case analysis)

Cox regression model	C index (95% CI)
PI-RADS model (n = 366)	
PRIMARY score	0.61 (0.49–0.72)
PI-RADS score	0.63 (0.54–0.72)
PI-RADS score + PRIMARY score	0.68 (0.58–0.77)
cT stage model (n = 960)	
PRIMARY score	0.58 (0.52–0.63)
cT stage	0.63 (0.57–0.68)
cT stage + PRIMARY score	0.65 (0.59–0.71)

CI = confidence interval; PI-RADS = Prostate Imaging-Reporting and Data System.

MARY. Optimism-corrected C index values were 0.60 for PI-RADS + PRIMARY, and 0.64 for cT stage + PRIMARY.

It has been shown that the PRIMARY score improves cancer diagnosis, and the score was incorporated into the PROMISE v2 framework for reporting PSMA PET to better characterize prostate findings with this imaging modality [5]. The PRIMARY score is a 5-point system comprising a combination of the intraprostate pattern (diffuse or focal), site (transition or peripheral zone), and intensity ($SUV_{max} \geq 12$) to overcome the limitations of benign PSMA activity, which is common within the prostate and reduces specificity. While PRIMARY scores of 1–4 focus on the pattern and site independent of intensity, PRIMARY score 5 is based on the PSMA intensity of the index lesion. It is known that greater PSMA intensity in the primary prostate tumor predicts both higher grade group and poorer outcomes following definitive primary therapy. PRIMARY score 5 thus provides information on tumor biology and aggressiveness on the basis of PSMA expression independent of other features such as prostatic pattern and site.

The OS prediction probability for csPC detected using the PRIMARY score alone was comparable to that using PI-RADS scores or clinical results without PET (cT stage). However, our results confirm the additive OS predictive value of the PRIMARY score when combined with either PI-RADS scores or cT stage.

Our study demonstrates for the first time the prognostic value of the PRIMARY score at initial staging of biopsy-confirmed PC. A PROMISE score of 5 was associated with shorter OS. We will continue to collect PRIMARY score data in our PROMISE registry (NCT06320223, promise-pet.org) for further analysis over longer OS follow-up.

Author contributions: Madeleine J. Karpinski had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Fendler, Hadaschik, Emmett.

Acquisition of data: Karpinski, Civan, Hoberück, Evangelista, Vondrak, Rasul, Forner, Güven, Ayati, Lanfranchi, Scholtissek, Miksch, Soeterik, Di Giorgio, Holzgreve.

Analysis and interpretation of data: Karpinski, Fendler, Emmett, Civan, Hoberück.

Drafting of the manuscript: Karpinski, Fendler, Emmett, Civan, Hoberück. **Critical revision of the manuscript for important intellectual content:** Karpinski, Hoberück, Fendler, Civan, Bundschuh, Thomas, Bjartell, Trägårdh, Soeterik, Evangelista, Vondrak, Rasul, Forner, Giorgio, Scholtissek, Miksch, Holzgreve, Lanfranchi, Rahbar, Hofman, Rauscher, Güven, Eiber, Ayati, Herrmann, Hadaschik, Emmett, Umutlu.

Statistical analysis: Karpinski.

Obtaining funding: Fendler, Herrmann.

Administrative, technical, or material support: Bundschuh, Thomas, Hofman, Herrmann, Rahbar, Rauscher, Eiber, Bjartell, Trägårdh, Hadaschik, Umutlu

Supervision: Fendler, Emmett.

Other: None.

Financial disclosures: Madeleine J. Karpinski certifies that all conflicts of interest, including specific financial interests and relationships and affiliations relevant to the subject matter or materials discussed in the manuscript (eg, employment/affiliation, grants or funding, consultancies, honoraria, stock ownership or options, expert testimony, royalties, or patents filed, received, or pending), are the following: Madeleine J. Karpinski reports ownership of stocks in Bayer, outside of the submitted work. Wolfgang P. Fendler reports research funding from SOFIE Bioscience, Bayer, Amgen, and AstraZeneca; consultant fees from Janssen, Perceptive, Bayer, Novartis, GE Healthcare, and Lilly; speaker fees from Janssen, Bayer, Novartis, Telix, GE Healthcare, Eczacıbaşı Monrol, ABX, Amgen, and UroTrials; and an image review role for Perceptive. Caner Civan reports a postdoctoral fellowship from Humboldt Foundation. Ralph Bundschuh reports consulting fees from Bayer Healthcare, Terumo, and Eisai; honoraria from Bayer Healthcare, Novartis, and Terumo; travel expenses from Blue Earth Therapeutics; advisory board participation for Eisai and Novartis; and medical writing assistance from Blue Earth Therapeutics; all outside of the submitted work. Anders Bjartell reports consulting, advisory board, or speaker fees from Accord, Astellas, AstraZeneca, Bayer, Ipsen, Johnson & Johnson, Pfizer, Sandoz, and Telix; institutional research grants from Astellas, AstraZeneca, Bayer, Johnson & Johnson, Movember, Roche, and Spectracure; co-founder/board member role for Glactone Pharma AB; and shares in Glactone Pharma AB, LIDDS AB, and WntResearch AB, all outside the submitted work. Laura Evangelista reports payment from GE Healthcare, Novartis, Ergea, and Menarini; reimbursement from Novartis; participation in advisory boards for Ergea; and an unpaid leadership role in EANM Women's Empowerment, all outside of the submitted work. Jonathan Miksch reports ownership of stocks in Bayer outside of the submitted work. Adrien Holzgreve reports grants and consulting fees from ABX; and leadership role in the Young German Society of Nuclear Medicine and Young Oncologists United, outside of the submitted work. Kambiz Rahbar reports consulting fees from Novartis, Bayer Healthcare, UroTrials, and Pharmtrace; honoraria or payments from Novartis, Bayer Healthcare, and ABX; and travel support from Bayer Healthcare and ABX, outside of the submitted work. Michael S. Hofman reports research support from the Australian Cancer Research Foundation, the Prostate Cancer Foundation, the National Health and Medical Research Council, and the Peter MacCallum Foundation; institutional research support and/or advisory board fees from AdvanCell, ANSTO, Bayer, Isotopia, Novartis, and MIM; personal consulting fees from Janssen, MSD, and Sanofi; and potential personal equity from AdvanCell. Isabel Rauscher reports consultant fees and research funding from Blue Earth Diagnostics, outside of the submitted work. Matthias Eiber reports consultant fees from Blue Earth Diagnostics, Novartis/AAA, Telix, Bayer,

RayzeBio, Point Biopharma, and Janssen Pharmaceuticals; research funding from Blue Earth Diagnostics and Bayer; speaker fees from Novartis/AAA, Eckert-Ziegler, Urotrials, and Janssen Pharmaceuticals; an image review role for Parexel and Bioclinica; a patent application for rhPSMA; royalties from sales of POSLUMA; and advisory board and/or steering committee roles for Novartis, Blue Earth Diagnostics, and Telix, all outside of the submitted work. Ken Herrmann reports grants or contracts from Novartis and Sofie Biosciences; consulting fees from Advanced Accelerator Applications, Amgen, AstraZeneca, Bain Capital, Bayer, Boston Scientific, Convergent, Curium, Debiopharm, EcoR1, Fusion, GE Healthcare, Immedica, Isotopen Technologien München, Janssen, Merck, Molecular Partners, NVision, POINT Biopharma, Pfizer, Radiopharm Theranostics, Rhine Pharma, Siemens Healthineers, Sofie Biosciences, Telix, Theragnostics, and ymabs; payment or honoraria for educational events from PeerVoice; travel support from Janssen; participation in a data safety monitoring board or advisory board for Fusion and GE Healthcare; and ownership of stocks in Sofie Biosciences, Pharma15, NVision, Convergent, Aktis Oncology, and AdvanCell, outside of the submitted work. Boris Hadaschik reports consulting fees from Janssen, Bayer, ABX, Astellas, Merck, Amgen, MSD/Pfizer, Novartis, BMS, Monrol, Onkowitz, POINT Biopharma, Ipsen, AstraZeneca, Lightpoint Medical, Telix, and Accord Healthcare; travel support from AstraZeneca, BMS, Janssen, Bayer, and Ipsen; grants or contracts from Janssen, Deutsche Forschungsgesellschaft, Novartis, and BMS; and participation in data safety monitoring boards for Janssen and ABX. Louise Emmett reports research funding from Novartis, Clarity Pharma, and Telix; advisory board fees from Clarity Pharma and AdvanCell; a consultant role for GE Healthcare and Astellas; travel support from Astellas; a PCF Challenge Award from the Prostate Cancer Foundation; and grant funding from Movember, St. Vincent's Clinic Foundation, and St. Vincent's Curran Foundation. The remaining authors have nothing to disclose.

Funding/Support and role of the sponsor: This research was partly funded by German Research Foundation (Deutsche Forschungsgemeinschaft, DFG, grant FE1573/7-1 / 714209 and RA 3222/5-1 / 714210, project number 564651276), Prostate Cancer Foundation TACTICAL Award No 22TACT01, Thera4Care by the Innovative Health Initiative Joint Undertaking (IHI JU) under grant agreement No 101172788, ILLUMINATE by the Innovative Health Initiative Joint Undertaking (IHI JU) under grant agreement No 101172722, and AstraZeneca (Grant ID# 95663957).

Data sharing statement: Access to the summary of participant data can be approved on an individual basis by an independent review committee

on reasonable request after publication. Requests should be directed to the corresponding author.

Acknowledgments: We thank the patients, clinicians, and investigators from all participating sites for their contribution. We also acknowledge significant contributions to data acquisition from Lena M. Unterrainer, Gabriel T. Sheikh, Sophie C. Kunte, Lukas Späth, Linus Hempel, and Anikka Roluna, and all other investigators from the PROMISE Registry group. Adrien Holzgreve is funded by Deutsche Forschungsgemeinschaft (grant 545058105).

Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.eururo.2025.10.013>.

References

- [1] Cornford P, Tilki D, van den Bergh RCN, et al. EAU-EANM-ESTRO-ESUR-ISUP-SIOG guidelines on prostate cancer. Arnhem, The Netherlands: European Association of Urology; 2025.
- [2] Kesch C, Vinsensia M, Radtke JP, et al. Intra-individual comparison of ¹⁸F-PSMA-1007 PET/CT, multiparametric MRI, and radical prostatectomy specimens in patients with primary prostate cancer: a retrospective, proof-of-concept study. *J Nucl Med* 2017;58:1805–10.
- [3] Scheltema MJ, Chang JJ, Stricker PD, et al. Diagnostic accuracy of ⁶⁸Ga-prostate-specific membrane antigen (PSMA) positron-emission tomography (PET) and multiparametric (mp)MRI to detect intermediate-grade intra-prostatic prostate cancer using whole-mount pathology: impact of the addition of ⁶⁸Ga-PSMA PET to mpMRI. *BJU Int* 2019;124(Suppl 1):42–9.
- [4] Emmett L, Papa N, Buteau J, et al. The PRIMARY score: using intraprostatic ⁶⁸Ga-PSMA PET/CT patterns to optimize prostate cancer diagnosis. *J Nucl Med* 2022;63:1644–50.
- [5] Seifert R, Emmett L, Rowe SP, et al. Second version of the Prostate Cancer Molecular Imaging Standardized Evaluation framework including response evaluation for clinical trials (PROMISE V2). *Eur Urol* 2023;83:405–12.
- [6] Guo S, Kang F, Ma S, et al. The PRIMARY score: diagnostic performance and added value compared with MRI in detecting clinically significant prostate cancer. *Clin Nucl Med* 2024;49:37–44.
- [7] Karpinski MJ, Rahbar K, Bogemann M, et al. Updated prostate cancer risk groups by prostate-specific membrane antigen positron emission tomography prostate cancer molecular imaging standardized evaluation (PPP2): results from an international multicentre registry study. *Eur Urol* 2025;88:484–95.