

# Attitudes toward large language model-based Artificial Intelligence systems as an information source for shared decision-making in radiation oncology

Rebecca Moser<sup>1</sup>, Lena M. Buchecker<sup>1</sup>, Jana Nano<sup>1</sup>, Nina A. Mayr<sup>2</sup>, Sophie T. Behzadi<sup>1</sup>, Sophia Kiesl<sup>1</sup>, Sophie Maier<sup>1</sup>, Luisa Allwohn<sup>1</sup>, Jacqueline Lammert<sup>3,4,5</sup>, Lisa C. Adams<sup>6</sup>, Max Tschochohei<sup>4,7</sup>, Stephanie E. Combs<sup>1,5,8</sup>, Kai J. Borm<sup>\*,1,9</sup> 

<sup>1</sup>Department of Radiation Oncology, TUM University Hospital, Klinikum Rechts der Isar, School of Medicine and Health, Technical University of Munich, Munich 81675, Germany

<sup>2</sup>College of Human Medicine, Michigan State University, East Lansing, MI 4882, United States

<sup>3</sup>Department of Gynecology and Center for Hereditary Breast and Ovarian Cancer, Technical University of Munich (TUM), School of Medicine and Health, Klinikum rechts der Isar, TUM University Hospital, Munich 81675, Germany

<sup>4</sup>Institute of Artificial Intelligence and Informatics in Medicine (AIIM), TUM University Hospital, Technical University of Munich (TUM), Munich 81675, Germany

<sup>5</sup>German Cancer Consortium (DKTK), partner site Munich, a partnership between DKFZ and TUM University Hospital, Munich, 81675, Germany

<sup>6</sup>Department of Diagnostic and Interventional Radiology, School of Medicine and Health, Klinikum rechts der Isar, TUM University Hospital, Technical University of Munich, Munich 81675, Germany

<sup>7</sup>Google Deutschland GmbH, Munich 80636, Germany

<sup>8</sup>Institute of Innovative Radiotherapy (IRT), Department of Radiation Sciences (DRS), Helmholtz Zentrum München, Oberschleißheim 85764, Germany

<sup>9</sup>Bavarian Center for Cancer Research (BZKF), Technical University Munich, Munich 81675, Germany

\*Corresponding author: Department of Radiation Oncology, TUM School of Medicine and Health, Ismaninger Straße 22, Munich 81675, Germany ([kai.borm@mri.tum.de](mailto:kai.borm@mri.tum.de)).

## Abstract

**Background:** Implementing structured shared decision-making (SDM) requires high-quality, reliable patient information. In radiation oncology, patients often have limited knowledge and misconceptions about therapy and side effects, affecting their decision-making. Large language model-based AI systems (LLMs) may help by providing evidence-based information in accessible language, but successful implementation depends on the willingness of patients and health care professionals (HCPs) to adopt these technologies.


**Methods:** A survey was conducted among patients undergoing radiation therapy and HCPs between 03/2024 and 02/2025. Data was collected using structured electronic questionnaires (32 items for patients, 35 for HCPs). The survey assessed sociodemographic characteristics, the status of SDM in oncology, sources of information relevant to SDM, and current and anticipated LLM applications. Data were analyzed using descriptive statistics and logistic regression analysis.

**Results:** The internet was the prime information source for patients ( $n=400$ ). Regarding current use of LLMs, a large discrepancy between patients and HCPs ( $n=200$ ) was observed (18.2% vs 69.5%). Although 77% of HCPs believed that patients will rely on LLMs in the future, only 29.1% of patients agreed. Most patients (65.8%) stated that even as LLMs improve, they will continue to trust physicians more; 46% of HCPs shared this view. Only 16.5% of patients were convinced that LLMs provide all relevant data for SDM in cancer care. Familiarity with technology was the strongest predictor of LLM use among patients.

**Conclusion:** Only a minority of radiation oncology patients currently use LLMs, and many remain skeptical about their future role—contrasting with the more optimistic expectations of HCPs.

**Key words:** radiotherapy; large language models; shared decision-making; cancer care; artificial intelligence; ChatGPT.

## Graphical abstract



### Implications for Practice

The use of large language model-based AI systems (LLMs) among healthcare professionals (HCPs) in radiation oncology is approximately four times higher than among patients. While most HCPs anticipate increasing patient reliance on LLMs, few patients share this view, consistently expressing greater trust in physicians. Despite LLMs' potential to support shared decision-making in cancer care, a critical barrier remains: the disconnect between professional optimism and patient skepticism. This challenge is not solely technical; effective implementation must focus on building patient trust, clearly demonstrating benefits, and ensuring that LLMs empower rather than alienate those they are intended to support.

### Introduction

Shared decision-making (SDM) is a central component of patient-centered care, particularly in oncology. Due to the technical nature of radiation therapy (RT), patients often experience higher levels of fear compared to other cancer treatments.<sup>1,4</sup> Moreover, their limited prior knowledge of RT may contribute to increased skepticism regarding potential side effects. Consequently, patients express a strong desire to be actively involved in treatment decision-making.<sup>5,6</sup>

To address this knowledge gap, evidence-based information sources and decision aids are essential for supporting patients' participation in SDM. These tools are crucial for helping patients understand their treatment options and enabling informed engagement. Nevertheless, although they are essential for SDM, their use in clinical settings remains limited. However, the development of individualized decision aids for each

patient is a time-intensive process that requires continuous review and updates to ensure relevance and actuality, making them not always readily available.

As a result, patients frequently turn to alternative sources of information such as internet search engines, leading to unverified information without the necessary accuracy, evidence, and personalization. Artificial intelligence (AI) systems, particularly those based on large language models-based AI systems (LLMs) have the potential to serve as an informational resource for both patients and healthcare professionals (HCPs). When properly configured, LLMs can provide evidence-based, physician-verified information in clear, easily understandable language. Several studies have investigated the quality and accuracy of AI systems in medical contexts—with and without the use of Retrieval-Augmented Generation, a method that enhances responses by integrating relevant external information retrieved

from trusted sources before generating output.<sup>7-12</sup> While the quality and accuracy of LLMs are of utmost importance, the equally important aspects of patient acceptance and engagement are often overlooked.<sup>13</sup> For example, Reis et al. conducted an online survey to evaluate patients' acceptance of digital medical advice provided by AI, physicians, or AI supervised by physicians. Their findings indicated that participants were less willing to follow recommendations when AI was involved in their generation.<sup>14</sup>

The primary objective of our survey study was to assess the interest of radiation oncology patients in utilizing LLMs as a source of medical information. Secondary objectives included evaluating the perceived future potential of LLMs as an informational resource and decision-support tool for SDM among both patients and HCPs.

## Methods

### Study design

The study was conducted at the Department of Radiation Oncology, Technical University of Munich, Germany, from March 2024 to February 2025. This prospective, cross-sectional survey study was supported by the Bavarian Center for Cancer Research and approved by the local institutional review board (2024-89-S-SB). All participants provided their written consent to take part in the anonymous online survey.

### Questionnaire design

Two structured questionnaires were developed in collaboration with experts in oncology, radiation oncology, SDM, and palliative care through moderated group discussions with professionals. Patients were not involved in the questionnaire design. In addition, a non-systematic literature search was conducted before designing the questionnaire to identify relevant topics, expectations, and concerns related to the use of LLMs in SDM. The survey items were designed to capture attitudes toward LLMs specifically, rather than toward technological innovations more broadly. The patient questionnaire consisted of 32 questions covering four categories: sociodemographic data, status of SDM in cancer care, current information sources for SDM, and current and future use of LLMs. The HCP questionnaire consisted of 35 questions divided into four sections: socio-demographic data, current status of SDM, current and future use of LLMs among patients, and personal experience with LLMs. To avoid misunderstandings, the patient questionnaire makes it clear that it refers specifically to "physicians," rather than to HCPs in general. Both questionnaires are available in the [Supplemental Material](#) (Questionnaires 1 and 2).

### Participants

Participants were required to have sufficient proficiency in German and the physical and mental ability to complete a structured questionnaire. The patient survey was administered to eligible cancer patients treated at the Department of Radiation Oncology.

Out of 438 patients invited to participate, 400 completed the survey, resulting in a response rate of 91.3%. Recruitment was consecutive during patients' visits; this may have introduced selection bias depending on which patients presented on specific days.

The second online survey was administered to HCPs in radiation oncology via a link that was distributed within the

department of radiation oncology, a social media link and the German Society of Radiation Oncology (DEGRO, approximately 2000 email addresses). A total of 200 HCPs responded. The exact response rate cannot be determined due to the data collection method.

At the beginning of the questionnaire, all participants were provided with the following introductory statement to avoid ambiguity concerning LLMs: "AI refers to computer programs or machines that are designed to perform tasks that would typically require human intelligence. Such systems are capable of learning, adapting to different situations, and making decisions without being explicitly programmed. LLMs, like ChatGPT, utilize machine learning and AI to engage in human-like conversations and respond to questions or requests."

### Statistical analysis

For the analysis of the primary outcome, patients' interest in using LLMs for medical information, and the secondary outcomes, data from the online survey were analyzed descriptively and visualized using SPSS, Excel, and R. Continuous variables are reported as mean  $\pm$  SD for normally distributed data and as median [interquartile range (IQR)] for skewed data. Categorical variables are presented as total numbers with the corresponding percentage.

To identify factors associated with patients' interest in LLMs, we performed a multivariable logistic regression analysis. The Likert items on LLM interest (outcome) and SDM interest were dichotomized into "Disagreement" (responses "Does not apply," "Rather does not apply," or "Neutral") versus "Agreement" (responses "Rather applies" or "Applies"), in order to provide a clinically interpretable distinction between patients with and without relevant interest. The original five-point Likert scales for technical affinity, SDM interest, and LLM interest were relabeled as ordered factors with descriptive levels. For the two interest items, responses of "Does not apply," "Rather does not apply," or "Neutral" were grouped into "Disagreement," and responses of "Rather applies" or "Applies" were grouped into "Agreement," yielding simple binary indicators. Because collapsing categories may reduce variability, we conducted a sensitivity analysis using ordinal logistic regression with the original ordered Likert responses to confirm the robustness of the findings. "Neutral" responses were used as the reference category for variables of LLM interest, technology affinity, and SDM interest. The model included age, gender, education level, technical affinity, and interest in SDM as independent variables. Education was recoded into a three-level factor (low, secondary, tertiary) using predefined categories. There was no missing data. Statistical significance was defined as  $\alpha = .05$ .

## Results

### Participants

Out of the 400 cancer patients, 225 (56.3%) patients were female and 175 (43.7%) were male with a median age of 64.1 years (range 17-89 years). Most patients had a lower (37.0%,  $n = 148$ ) or upper (21.0%,  $n = 84$ ) secondary level as their highest educational qualification. Predominant diagnoses were breast cancer (23.0%,  $n = 92$ ), head and neck cancer (16.0%,  $n = 64$ ), and brain tumors (15.3%,  $n = 61$ ) (Table S1).

HCPs' had a median age of 40.5 years (range 20-77 years), with 53.5% ( $n=107$ ) females. 82% ( $n=164$ ) worked at a university hospital, mostly in radiation oncology or with a direct interdisciplinary connection to radiation oncology. Most participants were physicians (50.5%,  $n=101$ ) or nurses (24.0%,  $n=48$ ) (Table S2).

### Technical affinity and preferred information sources

As shown in Figure 1, technical affinity was self-assessed on a five-point Likert scale. Among patients, 55.5% ( $n=222$ ) either "tended to agree" or "completely agreed" with having technical affinity, compared to 85.5% ( $n=171$ ) of HCPs. Only a small fraction of HCPs expressed disagreement; patient responses were more broadly distributed. Regarding information sources for treatment decisions with multiple possible answers, patients reported primarily using the internet (26.1%,  $n=148$ ), followed by second opinions from other doctors (23.9%,  $n=136$ ) and friends and acquaintances (16.4%,  $n=93$ ) (Figure 1). In contrast, HCPs recommended mainly second opinions (29%,  $n=173$ ), followed by informed consent documents (19.3%,  $n=115$ ) and informational brochures from professional societies (16.6%,  $n=99$ ).

### Current LLM experience and interest in LLMs as an information source for SDM

Although LLMs are increasingly discussed as potential tools in healthcare, current experience with LLMs and attitudes toward their use varied considerably between patients and HCPs. As shown in Figure 1, 81.8% ( $n=327$ ) of patients reported no prior experience with LLMs, versus 30.5% ( $n=61$ ) of HCPs. This major difference suggests that despite growing attention to AI in healthcare, patients remained skeptical about the role of LLMs in SDM. Only 43% ( $n=174$ ) of patients supported the idea of physicians using LLMs to generate personalized information, and just 28.7% ( $n=115$ ) expressed personal interest in using LLMs themselves. These findings suggest that patient skepticism toward LLMs may be driven by underlying factors such as digital confidence or interest in collaborative decision-making. Multivariable logistic regression analysis (Table S3) revealed that technical affinity was the strongest independent predictor of expressing interest in LLM use. Participants with higher self-assessed technical affinity were 40% more likely to express interest in LLMs (OR=1.40, 95% CI: 1.15–1.70,  $P<.001$ ), whereas each additional year of age was associated with 2% lower odds of reporting greater interest in LLMs (OR=0.98, 95% CI: 0.97–1.00,  $P=.04$ ). This association was further reflected in the distribution of technical affinity by age (Figure S1, see online supplementary material for a color version of this figure), showing that older patients more frequently reported low technical affinity. An interaction term of age and technical affinity was found not significant. Additionally, a strong interest in SDM was associated with substantially greater LLM interest (OR=3.42, 95% CI: 1.35–8.58,  $P=.009$ ). Gender showed no significant effect (OR=0.94, 95% CI: 0.61–1.47,  $P=.79$ ). Similarly, education level was not significantly associated with LLM interest, although participants with tertiary education tended to have higher odds compared to those with low education (OR=1.41, 95% CI: 0.82–2.43,  $P=.22$ ). Sensitivity analyses of the ordinal logistic regression confirmed these trends (Table S4).

### Future perspectives on LLMs as information sources

Only 37.8% ( $n=151$ ) of patients believed consulting an LLM regarding medical decisions could be helpful in the future. In contrast, a significantly higher proportion of HCPs anticipated an increased role of LLMs: 70.5% ( $n=141$ ) of HCPs expected that patients will increasingly use LLMs as an information source (Figure 2A), and 75.5% ( $n=151$ ) of HCPs could envision utilizing LLMs themselves to support patient decision-making. Furthermore, 77% of HCPs ( $n=154$ ) believed patients will increasingly rely on LLMs, independent of physician recommendation. 65.8% ( $n=263$ ) of patients expected to continue trusting physicians over LLMs, compared to 46% ( $n=92$ ) of HCPs, with 30% ( $n=60$ ) expressing doubt that patients will maintain a stronger trust in physicians over LLMs (Figure 2B).

Despite the rapid development of LLMs, only 16.5% ( $n=66$ ) of patients believed that LLMs could provide all relevant information for SDM in radiation oncology. In comparison, 32% ( $n=64$ ) of HCPs considered this likely (Figure 2C).

The majority of patients (71.5%,  $n=286$ ) and HCPs (81.5%,  $n=163$ ) agreed that LLMs cannot fully replace the physician-patient interaction within the context of SDM (Figure 2D). These findings highlight that even with optimism from HCPs, significant patient doubts remain, especially regarding trust and the desire for human involvement.


There were no differences in responses across the various HCP subgroups. The detailed responses to all items are provided in the supplementary dataset (Figures S2-S33, see online supplementary material for a color version of these figures).

### Discussion


Our study highlights significant differences in the perceptions of radiation oncology patients versus HCPs in radiation oncology regarding the role of LLMs in SDM. While most HCPs foresee a growing integration of LLMs into clinical workflows and express optimism regarding their potential, patients remain more hesitant, exhibiting limited current usage, lower trust, and concerns about the replacement of direct physician interactions. These differences are crucial to consider for the future development and implementation of LLMs in oncology care.

LLMs have a unique potential to address individualized patient education, which is essential for enabling the process of SDM. Even though RT is a crucial part of oncology, the topic of SDM in radiation oncology has been underexplored, and only very few studies specifically address SDM in this field.<sup>15-20</sup> Whereas some components of SDM, such as patient empowerment and HCP education, can easily be transferred from other specialties, information sources and decision aids specific to radiation oncology are complex due to technical, physical, and biological considerations. Therefore, there is a strong need for efficient individualized patient education in radiation oncology, particularly because patients usually have little prior knowledge or contact before therapy.

The creation of LLMs for patient education is currently underway in several scientific and commercial projects.<sup>12,21-25</sup> LLMs can provide evidence-based, personalized information and patient decision aids for the SDM process in radiation oncology that are accessible, easy to understand, and reduce



**Figure 1.** Technological affinity, information behavior, and perspectives on AI in oncology between patients (A) and professionals (B). Technological affinity: “Self-assessment of technical affinity, Q: Would you consider yourself to be technically savvy?” Current experience with LLMs: Q: Have you ever interacted with an artificial intelligence (e.g., ChatGPT)?” Interest in future use of LLMs: Patients: Q: “I would generally be interested in asking questions to a specially trained chatbot.” Professionals: Q: “Do you think that patients will increasingly rely on AI and chatbots in the future, regardless of professional recommendations?” Information sources: Patients: Q: “Where did you obtain this additional information (beyond what was provided by your doctors)?” Professionals: Q: “What sources of information do you provide or recommend to patients for making informed decisions in oncology?” LLM, large language models.



**Figure 2.** Perspectives on the future use of LLMs in SDM in cancer care. (A) Patients: Q: “Do you think that consulting artificial intelligence could be helpful to you as a patient when making medical decisions?” Professionals: Q: “Could you imagine using AI more frequently in the future to better inform patients regarding oncological decisions?” (B) Patients: Q: “I would trust the statements of doctors more than those of artificial intelligence in the decision-making process.” Professionals: Q: “In the future, will patients trust the statements of doctors more than those of AI in the decision-making process?” (C) Patients: Q: “AI provides me with all the information relevant to the decision-making process.” Professionals: Q: “Does AI provide patients with all the information relevant to the decision-making process?” (D) Patients: Q: “Do you think that AI will replace doctor-patient conversations for shared decision-making in the future?” Professionals: Q: “Do you think that AI will replace doctor-patient conversations for shared decision-making in the future?” LLM, large language models; SDM, shared decision-making.

the need for patients to search confounding and often unverified sources on the internet.

Although patients often use the internet for [supplementary information](#), this does not imply trust in its reliability. HCPs recommend more structured, evidence-based sources, which patients perceive as more trustworthy. Therefore, frequent online information seeking can coexist with skepticism toward LLMs, emphasizing the need to demonstrate reliability and clinical value when integrating LLMs into SDM.

While HCPs see the potential of LLMs, our results show that most patients do not embrace this potential and express low interest in LLMs for their radiation oncology care. This skepticism may be driven by the current limitations of LLMs, such as the possibility of generating inaccurate or unverified information (“hallucinations”), which patients are less able to assess than professionals.<sup>21,22</sup>

While LLM-supported tools may eventually become embedded in routine healthcare workflows irrespective of individual interest, their meaningful use within SDM continues to depend on patient engagement and perceived autonomy. Our participants did not explicitly express concerns that LLM-based applications might be imposed on them in future care scenarios; however, the comparatively low intention among patients to rely on LLMs suggests a desire to retain control over how such technologies are used in decision-making processes. In contrast, HCPs—who reported both greater familiarity and higher expectations for future LLM utility—may anticipate a more rapid integration of these systems into clinical practice. This divergence underscores that even if LLMs become ubiquitous, acceptance cannot be assumed, particularly in contexts such as SDM, where trust, interpersonal communication, and

patient empowerment remain central. Ensuring that LLM use supports rather than replaces patient autonomy will therefore be critical as implementation progresses. Moreover, our group undertakes a systematic analysis to delineate the circumstances in which the deployment of AI and LLMs requires the explicit procurement of patient consent from the patients’ perspective.

The gap between patient and HCP expectations may also be influenced by age and technical affinity. Older patients more frequently reported low technical affinity, which was the strongest predictor of LLM interest. However, age alone was not the primary factor even though the surveyed professionals had a median age of 40, while patients had a median age of 64; interest in SDM and comfort with technology appear more important. Patients with strong SDM preferences were markedly more interested in using LLMs, indicating that those who seek active involvement in decisions may view such tools as supportive rather than substitutive. Gender, education, and other demographic properties did not significantly influence LLM acceptance.

The most important impact factor for LLMs skepticism was low technical affinity. Although current skepticism toward LLMs is largely driven by concerns regarding the insufficient quality and credibility of LLMs in comparison to human experts, the rapid progress in both scientific and commercial projects as well as increasing integration of electronic communication in healthcare suggests that such concerns may become less justified in the future. This underscores the potential of LLMs to improve patient–clinician interactions.<sup>23</sup> However, recent population-based studies show that LLM use is already widespread and increasing, both for general and health-related

purposes. This suggests that the comparatively low use observed in our cohort may be related to the specific characteristics of oncology patients rather than general public trends.<sup>24,25</sup>

While familiarity with LLMs is expected to increase over time—as is typically the case with technological innovations—experience suggests that the acceptance of such tools will likewise grow. This development can be supported by continuously promoting education and exposure to the topic, as well as by providing guidance to patients in their use.

The discrepancy between patients' and HCPs' perspectives may additionally reflect differences in professional exposure: HCPs are generally more familiar with digital innovations and may therefore anticipate their potential benefits more readily, whereas patients may experience LLMs as abstract, complex, or potentially risky. Such psychological barriers—such as uncertainty, fear, or a perceived lack of control—were not assessed in our study but represent important factors for future research.

The application of LLMs in the preparatory phase of medical consultations may transform the dynamics of patient–physician communication by establishing a shared foundation of knowledge. Such an approach may enable patients to perceive the interaction as more equitable and to prepare targeted questions in advance, ultimately fostering patient empowerment and supporting SDM. However, a potential drawback lies in the risk that patients may be exposed to unfiltered or distressing medical information that they are unable to interpret or emotionally process without professional support. In such cases—particularly when discussing sensitive topics such as prognostic outcomes in metastatic cancer—human interaction and empathetic communication remain indispensable.<sup>26</sup> This may also account for the continued higher level of trust patients place in physicians, based on the assumption that physicians demonstrate empathy and draw upon their clinical experience during the consultation. These considerations highlight why, despite ongoing advancements in LLM technologies, in our study three-quarters of both patients and HCPs did not believe that LLMs would be able to fully replace physician–patient interactions and believe that LLMs should function as supportive tools rather than autonomous decision-makers within SDM. For this reason, the Technical University of Munich is currently preparing a randomized trial in which different application methods of LLMs as information sources for patients in the SDM process will be investigated.

Structured integration strategies may help strengthen trust and demonstrate clinical value—for example, supervised use of LLMs before consultations or incorporating LLM-generated information into patient education workflows. Such models could help patients acquire preliminary knowledge, bridge waiting times efficiently, and bring targeted questions into the clinical encounter, while ensuring that any unclear or anxiety-provoking content is directly discussed with their physician.

### Limitations

In our study, the patient sample was recruited from a single radiation oncology department in Germany and therefore represents a small and specific patient population with severe illnesses; results may differ in other cultural or healthcare contexts.<sup>27</sup> Recruitment occurred consecutively during clinic visits, which may have introduced a selection bias depending

on which patients were present on specific days. HCPs participated via online link, likely leading to overrepresentation of digitally engaged individuals and inflating positive attitudes toward LLMs. Survey-based research inherently includes self-selection and social desirability bias. Although we do not assume substantial changes in understanding or awareness of LLMs during the one-year recruitment period, temporal effects cannot be completely ruled out. We asked participants about potential confounders such as digital experience, technical affinity, and education, which are reported in the results and [supplemental material](#). However, disease-specific factors, emotional burden, and psychological variables were not captured. Finally, challenges such as data privacy, reliability of AI outputs, workflow integration, and regulatory considerations represent additional barriers to successful implementation and should be addressed in future studies.

The current study results clearly indicate that the successful implementation of LLMs in clinical practice depends not only on technological development but also on systematic education and training for both patients and HCPs. Additional barriers—such as technical reliability, privacy protection, patient confidentiality, regulatory constraints, and the practical integration into clinical workflows—must be addressed. Only by meeting these requirements LLMs can unfold their full potential as safe, supportive, and meaningful tools within modern radiation oncology care.

To the best of our knowledge, our study is the first to compare the perspectives of patients and HCPs regarding the use of LLMs in SDM in radiation oncology, thereby addressing an existing knowledge gap with high impact on the success and adoption of LLMs. While the potential of SDM is currently the focus of extensive research, providing structured guidance and education and preserving physician–patient interactions as the core of SDM are essential for its future applicability. Future research should explore longitudinal changes in patient attitudes, the impact of exposure and education on acceptance, and strategies for integrating LLMs into routine clinical workflows safely and effectively.

### Conclusion

LLMs hold significant promise in supporting SDM in cancer care. While HCPs involved in RT generally maintain a positive outlook on the potential of LLMs, patients remain more skeptical and show lower interest in using LLMs. The key challenges will not only involve advancing and tailoring LLMs specifically for radiation oncology but also carefully exploring how to integrate these tools into clinical practice in a way that prioritizes patient interests and ensures tangible benefits for patients.

### Author contributions

Rebecca Moser (Conceptualization, Data curation, Formal analysis, Visualization, Writing—review & editing), Lena Marie Buchecker (Formal analysis, Supervision, Validation, Visualization, Writing—review & editing), Jana Nano (Supervision, Writing—review & editing), Nina A. Mayr (Writing—review & editing), Sophie T. Behzadi (Writing—review & editing), Sophia Kiesl (Writing—review & editing), Sophie Maier (Writing—review & editing), Luisa

Allwohn (Writing—review & editing), Jacqueline Lammert (Writing—review & editing), Lisa Christine Adams (Writing—review & editing), Max Tschochohei (Writing—review & editing), and Stephanie E. Combs (Conceptualization, Data curation, Funding acquisition, Supervision, Validation, Writing—original draft, Writing—review & editing)

## Supplementary material

Supplementary material is available at *The Oncologist* online.

## Funding

This research has received support from the Bavarian Center for Cancer Research (BZKF) in 2024 (BZKF-Young Scientist Fellowship).

## Conflicts of interest

The authors have no relevant financial or non-financial interests to disclose.

## Data availability

The datasets analyzed during the current study are available from the corresponding author upon reasonable request.

## Ethics approval and consent to participate

The study was approved by the local institutional review board (2024-89-S-SB), and all participants provided written informed consent. The procedures set out in this trial, pertaining to the conduct, evaluation, and documentation of this trial, are designed to ensure that all persons involved in the trial by Good Clinical Practice and the ethical principles described in the applicable version of the Declaration of Helsinki (2008 Version of the Declaration of Helsinki, adopted at the 59th WMA General Assembly, Seoul, October 2008). The trial will be carried out in keeping with local legal and regulatory requirements.

## References

- Mose S, Budischewski KM, Rahn AN, Zander-Heinz AC, Bormeth S, Böttcher HD. Influence of irradiation on therapy-associated psychological distress in breast carcinoma patients. *Int J Radiat Oncol Biol Phys.* 2001;51:1328-1335.
- Halkett GK, Kristjanson LJ, Lobb EA. 'If we get too close to your bones they'll go brittle': women's initial fears about radiotherapy for early breast cancer. *Psychooncology.* 2008;17:877-884.
- Piroth MD, Draia S, Jawad JA, Piefke M. Anxiety and depression in patients with breast cancer undergoing radiotherapy: the role of intelligence, life history, and social support—preliminary results from a monocentric analysis. *Strahlenther Onkol.* 2022;198:388-396.
- Lewis F, Merckaert I, Liénard A, et al. Anxiety and its time courses during radiotherapy for non-metastatic breast cancer: a longitudinal study. *Radiother Oncol.* 2014;111:276-280.
- Gaston CM, Mitchell G. Information giving and decision-making in patients with advanced cancer: a systematic review. *Soc Sci Med.* 2005;61:2252-2264.
- Chewning B, Bylund CL, Shah B, Arora NK, Gueguen JA, Makoul G. Patient preferences for shared decisions: a systematic review. *Patient Educ Couns.* 2012;86:9-18.
- Chen S, Kann BH, Foote MB, et al. Use of artificial intelligence chatbots for cancer treatment information. *JAMA Oncol.* 2023;9:1459-1462.
- Kuang YR, Zou MX, Niu HQ, Zheng BY, Zhang TL, Zheng BW. ChatGPT encounters multiple opportunities and challenges in neurosurgery. *Int J Surg.* 2023;109:2886-2891.
- Kim JK, Chua M, Rickard M, Lorenzo A. ChatGPT and large language model (LLM) chatbots: the current state of acceptability and a proposal for guidelines on utilization in academic medicine. *J Pediatr Urol.* 2023;19:598-604.
- Rajjoub R, Arroyave JS, Zaidat B, et al. ChatGPT and its role in the decision-making for the diagnosis and treatment of lumbar spinal stenosis: a comparative analysis and narrative review. *Global Spine J.* 2024;14:998-1017.
- Yeo YH, Samaan JS, Ng WH, et al. Assessing the performance of ChatGPT in answering questions regarding cirrhosis and hepatocellular carcinoma. *Clin Mol Hepatol.* 2023;29:721-732.
- G Xiong, Q Jin, Z Lu, A Zhang, eds. *Benchmarking Retrieval-Augmented Generation for Medicine.* Association for Computational Linguistics; 2024.
- Choudhury A, Shamszade H. Investigating the impact of user trust on the adoption and use of ChatGPT: survey analysis. *J Med Internet Res.* 2023; 25: e47184.
- Reis M, Reis F, Kunde W. Influence of believed AI involvement on the perception of digital medical advice. *Nat Med.* 2024;30:3098-3100.
- Leech M, Katz MS, Kazmierska J, et al. Empowering patients in decision-making in radiation oncology - can we do better? *Mol Oncol.* 2020;14:1442-60.
- Shabason JE, Mao JJ, Frankel ES, et al. Shared decision-making and patient control in radiation oncology: Implications for patient satisfaction. *Cancer.* 2014;120:1863-70.
- Shakespeare TP, Westhuyzen J, Lim Yew Fai T, et al. Choosing between conventional and hypofractionated prostate cancer radiation therapy: Results from a study of shared decision-making. *Rep Pract Oncol Radiother.* 2020;25:193-9.
- Smith SK, Nathan D, Taylor J, et al. Patients' experience of decisionmaking and receiving information during radiation therapy: A qualitative study. *Eur J Oncol Nurs.* 2017;30:97-106.
- Wang EH, Gross CP, Tilburt JC, et al. Shared decision making and use of decision AIDS for localized prostate cancer: perceptions from radiation oncologists and urologists. *JAMA Intern Med.* 2015; 175:792-9.
- Woodhouse KD, Tremont K, Vachani A, et al. A Review of Shared Decision-Making and Patient Decision Aids in Radiation Oncology. *J Cancer Educ.* 2017;32:238-45.
- Adams LC, Truhn D, Busch F, et al. Leveraging GPT-4 for post hoc transformation of free-text radiology reports into structured reporting: a multilingual feasibility study. *Radiology.* 2023;307:e230725.
- Chen Z, Hernández Cano A, Romanou A, et al. Scaling Medical Pretraining for Large Language Models, CoRR, abs/2311.16079, 2023. <https://doi.org/10.48550/arXiv.2311.16079>
- Jiang LY, Liu XC, Nejatian NP, et al. Health system-scale language models are all-purpose prediction engines. *Nature.* 2023;619:357-362.
- Y Labrak, A Bazoge, E Morin, P-A Gourraud, M Rouvier, R Dufour, eds. *BioMistral: A Collection of Open-Source Pretrained Large Language Models for Medical Domains.* Association for Computational Linguistics; 2024.
- Yang X, Chen A, PourNejatian N, et al. A large language model for electronic health records. *NPJ Digit Med.* 2022;5:194.
- Mittelstadt B, Wachter S, Russell C. To protect science, we must use LLMs as zero-shot translators. *Nat Hum Behav.* 2023;7:1830-1832.

27. Huang L, Yu W, Ma W, et al. A survey on hallucination in large language models: principles, taxonomy, challenges, and open questions. *ACM Trans Inf Syst.* 2025;43:1-55.
28. Benary M, Wang XD, Schmidt M, et al. Leveraging large language models for decision support in personalized oncology. *JAMA Netw Open.* 2023;6:e2343689.
29. Chatterji A, Cunningham T, Deming DJ, et al. How People Use ChatGPT. NBER Working Paper 34255; 2025.
30. Kacperski C, Ulloa R, Bonnay D, Kulshrestha J, Selb P, Spitz A. Characteristics of ChatGPT users from Germany: implications for the digital divide from web tracking data. *PLoS One.* 2025;20:e0309047.
31. Cè M, Chiarpello V, Bubba A, et al. Exploring the role of ChatGPT in oncology: providing information and support for cancer patients. *BioMedInformatics.* 2024;4:877-888.
32. Hussain S, Alherz M, Albazee E, et al. Investigating public perception on use of ChatGPT in initial consultations prior to healthcare provider consultations. *Annals of Medicine and Surgery.* 2024;9900: <https://doi.org/10.1097/MS9.0000000000002697>