JAMA Oncology | Original Investigation

# Consolidative Thoracic Radiotherapy With Atezolizumab Maintenance in Extensive-Stage Small Cell Lung Cancer
## The Phase 2 TREASURE Randomized Clinical Trial (AIO-TRK-0320)

Farastuk Bozorgmehr, MD; Inn Chung, PhD; Rouven Behnisch, PhD; Fabian Weykamp, MD; Sebastian Adeberg, MD; Tobias Overbeck, MD; Rami El Shafie, MD; Nadia Maguire, MD; Stefan A. Koerber, MD; Eva Lotte Buchmeier, MD; Michaela Hammer-Hellmig, MD; Thomas Gauler, MD; Christoph Pöttgen, MD; Martin Wermke, MD; Esther G. C. Troost, MD; Konrad Kokowski, MD; Barbara Röper, MD; Anke Reinacher-Schick, MD; Nicole-Sophie Consdorf, MD; Matthias Ulmer, MD; Arndt-Christian Müller, MD; Thomas Wehler, MD; Silla Hey-Koch, MD; Jürgen Alt, MD; Marcus Stockinger, MD; Jonas Kuon, MD; Marc Bischof, MD; Bernd Lamprecht, MD; Hans Geinitz, MD; Christian Lerchenmüller, MD; Jan Kriz, MD; Markus Rauter, MD; Wolfgang Raunik, MD; Bernd Schmidt, MD; Andrej Stupavsky, MD; Cornelia Kropf-Sanchen, MD; Gerlinde Schmidtke-Schrezenmeier, MD; Thomas Wiegel, MD; Amanda Tufman, MD; Farkhad Manapov, MD; Marie Merling, MSc; Claus Peter Heussel, MD; Markus Polke, MD; Martin Reck, MD; Salah-Eddin Al-Batran, MD; Michael Thomas, MD; Petros Christopoulos, MD; Stefan Rieken, MD

**IMPORTANCE** Chemoimmunotherapy followed by immunotherapy maintenance is the standard first-line treatment for extensive-stage small cell lung cancer (ES-SCLC), yet data regarding efficacy and safety of consolidative thoracic radiotherapy (TRT) are lacking.

**OBJECTIVE** To determine whether combining consolidative TRT with immunotherapy maintenance in ES-SCLC is safe and improves patients' overall survival (OS) and progression-free survival (PFS).

**DESIGN, SETTINGS, AND PARTICIPANTS** This was a multicenter open-label phase 2 randomized clinical trial recruiting patients from September 2020 to August 2022 in Germany and Austria, with the last follow-up visit in September 2024. Eligible patients had ES-SCLC with at least stable disease after induction chemoimmunotherapy (carboplatin–etoposide–atezolizumab). Although the database was locked in April 2025, a post hoc survival update was performed in April 2026. Data were analyzed from April 2025 to April 2026.

**INTERVENTIONS** Randomized (1:1) to either atezolizumab maintenance combined with consolidative TRT (30 Gy in 10 fractions) (arm A) or atezolizumab maintenance alone (arm B).

**MAIN OUTCOME AND MEASURE** OS (time from randomization to death due to any cause).

**RESULTS** Of 96 patients assessed for eligibility, 68 patients were randomized; recruitment was prematurely terminated due to safety concerns. Median OS in the combination arm was numerically shorter compared to atezolizumab only (6.7 months [95% CI, 5.1-9.0] vs 13.4 months [95% CI, 10.7-17.5]; HR, 1.55 [95% CI, 0.90-2.69]; $P = .34$), while median PFS was similar (2.4 months [95% CI, 1.3-3.9] vs 2.6 months [95% CI, 1.2-3.9]; HR, 0.92 [95% CI, 0.54-1.55]; $P = .85$). TRT plus atezolizumab was accompanied by higher frequency of severe adverse events (SAEs) (19 patients [61.3%] vs 6 patients [18.2%]; $P < .001$) and fatal outcomes (6 patients [19.4%] vs 1 patient [3.0%]; $P = .04$). Safety analysis revealed predominance of infection and respiratory disorder–related SAEs, possibly facilitated by a depletion of lymphocytes observed specifically in patients after TRT, and by a lower single breath diffuse capacity of the lungs for carbon monoxide in patients with fatal AEs in arm A. No further risk factors were identified, given that baseline characteristics were well balanced between both arms and between patients with and without SAEs, including fatal SAEs.

**CONCLUSIONS AND RELEVANCE** In this randomized clinical trial, TRT combined with immunotherapy increased toxic effects in unselected patients with ES-SCLC without survival benefit, presumably due to radiation-induced lymphocyte depletion facilitating infections. Cautious patient selection and further investigation to identify those who may benefit without undue risk are required.

**TRIAL REGISTRATION** ClinicalTrials.gov Identifier: NCT04462276

*JAMA Oncol*. 2026;12(9):965-972. doi:10.1001/jamaoncol.2026.2330
Published online July 9, 2026.

Visual Abstract
Editor's Note page 973
Supplemental content

**Author Affiliations:** Author affiliations are listed at the end of this article.

**Corresponding Author:** Farastuk Bozorgmehr, MD, Department of Thoracic Oncology, Thoraxklinik Heidelberg gGmbH, Röntgenstrasse 1, 69126 Heidelberg, Germany (farastuk.bozorgmehr@med.uni-heidelberg.de).



Downloaded from jamanetwork.com by DKTZ-DT Krebsforschungszentrum user on 11/13/2026

Atezolizumab combined with carboplatin–etoposide was established as the new standard of care for extensive-stage small cell lung cancer (ES-SCLC) based on the IMpower133 randomized clinical trial,[1] which reported a significant overall survival (OS) benefit in the atezolizumab arm compared to placebo. Combining photon radiotherapy (RT) with immunotherapy is being explored as a synergistic approach, based on the hypothesis that RT itself promotes an in situ vaccination through tumor-associated antigen exposure and immune cells recruitment.[2] In non–small cell lung cancer, this combination has shown encouraging improvements in outcomes, with good tolerability.[3-5] For ES-SCLC, the role of consolidative thoracic radiotherapy (TRT) added to immunotherapy remains poorly defined.[6-8] The CREST[9] trial previously demonstrated a modest OS benefit with TRT after chemotherapy in the preimmunotherapy era. The TREASURE trial aimed to further improve outcomes demonstrated by the CREST and IMpower133 results by adding TRT to atezolizumab maintenance for treatment of patients with ES-SCLC.

### Key Points

**Question** Is consolidative thoracic radiotherapy (TRT) added to standard immunotherapy maintenance in first-line extensive-stage small cell lung cancer (ES-SCLC) safe and more effective than immunotherapy alone?

**Findings** This phase 2 randomized clinical trial found that TRT plus immunotherapy unexpectedly resulted in significantly more frequent and severe toxic effects, particularly infections, without improving survival; therefore, recruitment was halted prematurely. Persistent lymphocyte depletion after TRT suggested a potential contributor to the observed toxic effects.

**Meaning** These findings indicate that in unselected patients with ES-SCLC, consolidative TRT combined with immunotherapy maintenance increased toxic effects without efficacy benefit, arguing against routine use other than in carefully selected patients; further investigation of radiation-induced lymphocyte depletion as a potential risk factor is warranted.

## Methods

This study was approved by the ethics committees at University of Heidelberg (AFmu-199/2020; 06/2020) and Johannes Kepler University of Linz (ECS 1097/2021; 05/2021). Written informed consent was obtained from all participants. The TREASURE randomized clinical trial (AIO-TRK-0320)[10] protocol is available in Supplement 1, and the statistical analysis plan, in Supplement 2. The Consolidated Standards of Reporting Trials (CONSORT) reporting guidelines were followed.

### Study Design and Sample Size

This multicenter phase 2 randomized clinical trial included patients with ES-SCLC and at least stable disease after induction with carboplatin–etoposide–atezolizumab at 20 sites in Germany and Austria. Sample size calculation was based on the primary end point OS (2-sided log-rank; $\alpha = .05$), assuming 20% absolute improvement in 12-month survival rate (72% vs 52%; hazard ratio [HR], 0.57), requiring 66 events and 92 patients (46 per arm) for 80% power. Including dropouts, 104 randomized patients were planned, and recruitment began in September 2020. Due to increased incidence of fatal SAEs in arm A vs arm B, the SMC recommended halting recruitment after enrollment of 68 participants (34 patients in each arm). Hence, statistical power was drastically reduced, and interpretation of results remains descriptive.

### Treatment Arms

Patients were randomized 1:1 to atezolizumab maintenance combined with consolidative TRT (30 Gy in 10 fractions) (arm A) or atezolizumab maintenance alone (arm B), stratified by presence of brain metastases, response after induction therapy, and prophylactic cranial irradiation following allocation by variance minimization procedure. Target volume delineation was based on postinduction volume of thoracic primary tumor and lymph node metastasis (dosimetric details are available in eTable 1 in Supplement 3).

### Outcomes

Primary end point was OS, defined as time from randomization to death due to any cause. Secondary end points included progression-free survival (PFS) and frequency and severity of adverse events (AEs) and serious adverse events (SAEs).

### Safety Monitoring

A predefined safety interim analysis assessed grade 3 or greater pneumonitis events after enrollment of 23 patients in arm A, obliging a recruitment halt if more than 1 event occurred, while a Safety Monitoring Committee (SMC) oversaw study progress.[10] After a fatal pneumonitis incident in a patient with preexisting interstitial lung disease (exclusion criterion), mandatory central review for preexisting interstitial lung disease was implemented in April 2021. The last patient was included in August 2022.[11] Treatment and follow-up continued until September 2024. Database lock occurred in April 2025. A post hoc OS analysis was based on an update in April 2026.

### Statistical Analysis

Primary and secondary efficacy analysis used stratified log-rank tests and multivariable Cox regression in the intention-to-treat (ITT) population ($\alpha = .05$) with point estimates of the HR and 95% profile likelihood confidence intervals. Sensitivity analyses were performed in the per-protocol and adjusted ITT set (ITT without fatal adverse events [AEs]).


Safety analysis was conducted in the safety population according to actual treatment in all patients who received at least 1 dose of atezolizumab (both arms) and started TRT (arm A) (eFigure 1 and the eMethods in Supplement 3).

Descriptive statistics, $\chi^2$ tests, *t* tests, and 2-proportion *Z*-tests were used to compare baseline characteristics and safety data between arms. Statistical tests were 2-tailed and *P* values < .05 were considered statistically significant. Data were analyzed from April 2025 to April 2026 using SAS, version 9.4, TS Level 1M6 (SAS Institute Inc).



Downloaded from jamanetwork.com by DKTZ-DT Krebsforschungszentrum user on 11/13/2026

Figure 1. Kaplan-Meier Estimates of Overall Survival (OS) and Progression-Free Survival (PFS) in the Intention-to-Treat Population




Median OS in arm A was shorter than arm B, without statistical significance (6.7 [95% CI, 5.1-9.0] months vs 13.4 [95% CI, 10.7-17.5] months; HR, 1.55 [95% CI, 0.90-2.69]; *P* = .34). Median PFS was similar (2.4 months [95% CI, 1.3-3.9] vs 2.6 months [95% CI, 1.2-3.9]; HR, 0.92 [95% CI, 0.54-1.55]; *P* = .85).

Stratification factors included in the stratified *P* value and Cox model were the same as those used for randomization. HR indicates hazard ratio; TRT, consolidative thoracic radiotherapy.

## Results

Of 96 patients screened, 68 patients (71%) were randomized (eFigure 1 in Supplement 3). The last patient was followed up until September 2024 per protocol.

Median OS in arm A was shorter compared to arm B, without statistical significance (6.7 [95% CI, 5.1-9.0] months vs 13.4 [95% CI, 10.7-17.5] months; HR, 1.55 [95% CI, 0.90-2.69]; *P* = .34) (**Figure 1**A). Estimated 1-year and 2-year OS rates in arm A were 30.3% (95% CI, 15.9%-46.1%) and 24.2% (95% CI, 11.4%-39.6%), and in arm B, 56.6% (95% CI, 37.9%-71.6%) and 15.7% (95% CI, 5.7%-30.2%), respectively. Post hoc OS analysis after additional 19 months of follow-up revealed similar results (eFigure 2 in Supplement 3). PFS was similar between arms (median, 2.4 [95% CI, 1.3-3.9] months vs 2.6 [95% CI, 1.2-3.9] months; HR, 0.92 [95% CI, 0.54-1.55]; *P* = .85) (Figure 1B). Efficacy outcomes and baseline parameters were consistent across adjusted ITT, per-protocol, and safety population (eFigure 2 and eTables 2 and 3 in Supplement 3).

Among the first 23 patients in arm A, only 1 case of grade 3 or greater pneumonitis occurred, fulfilling interim safety criteria. However, when the SMC noted increased fatal nonpneumonitis SAEs in arm A, recruitment was paused (August 2022), and then permanently stopped in December 2022 when an unplanned preliminary OS analysis (shared exclusively with the SMC to ensure study integrity) revealed a disadvantage of the combination therapy in arm A.[11]

In the final safety analysis, 30 patients (96.8%) in arm A and 25 (75.8%) in arm B experienced toxic effects (95% CI, 3.6%-39.5%; *P* = .02) (eTable 4 in Supplement 3). SAEs occurred in 19 patients in arm A (61.3%) vs 6 in arm B (18.2%; 95% CI, 16.5%-63.6%; *P* < .001). Treatment-related AEs (trAEs) and SAEs (trSAEs) were more frequent in arm A (trAEs, 71.0% vs 30.3%; *P* = .001; trSAEs, 29.0% vs 6.1%; *P* = .01), with greater differences for grade 3 or higher (eFigure 3 and eTable 4 in Supplement 3). Fatal AEs were significantly more frequent in arm A compared with arm B (6 patients [19.4%] vs 1 patient [3.0%]; *P* = .04). Causal attribution diverged between assessors (eTable 5 in Supplement 3); while the treating investigators considered 4 fatalities in arm A as unrelated to either atezolizumab or TRT, the SMC subsequently classified 3 of these as possibly or probably related to study treatment. Thus, causal relationship was not clearcut on a case-by-case basis.

Patients receiving TRT were more prone to microbial infections (urinary and pulmonary) and gastrointestinal (dysphagia and esophagitis), respiratory (particularly pneumonitis), fever, and skin AEs (**Figure 2**; eTable 6 in Supplement 3).


No baseline differences between arms accounted for the higher (S)AE incidence in arm A, including Eastern Cooperative Oncology Group Performance Status, Charlson Comorbidity Index, pulmonary function test results, thoracic tumor burden (**Table 1**) or tumor location (eTable 7 in Supplement 3), and neither clinical nor dosimetric characteristics were associated with SAE occurrences in arm A (eTables 8 and 9 in Supplement 3). However, the mean (SD) of baseline diffuse capacity of the lungs for carbon monoxide single breath (DLCO SB) was significantly worse in patients with fatal AEs vs other patients in arm A (45.3 [13.7] vs 56.7 [11.4]; *P* = .04; eTable 8 in Supplement 3), and organ-at-risk (OAR) analysis revealed a statistically significant, but clinically most likely not relevant, modest increase of total body dose exposure for them (median [range], 4.7 [3.0-6.6] Gy vs 3.4 [1.7-7.0] Gy; *P* = .04; eTable 9 in Supplement 3). Gross target volume was numerically larger in patients with fatal AEs (median [range], 88.6 [12.4-706.3] mL vs 45 [0-301.0] mL; *P* = .08). Radiation doses and volumes were within typical clinical ranges and below established OAR thresholds (eTable 1 in Supplement 3).

In a time-to-event analysis, TRT correlated positively (HR, 2.47; 95% CI, 1.15-5.32; *P* = .01), while atezolizumab expo-



Downloaded from jamanetwork.com by DKTZ-DT Krebsforschungszentrum user on 11/13/2026

Figure 2. Categories of (Serious) Adverse Events in Arm A (Atezolizumab Maintenance With TRT) vs Arm B (Atezolizumab Maintenance Only)




Adverse event categories, as defined by the Medical Dictionary for Regulatory Activities, which occurred at least 3 times are reported with the number of occurrences and exact 95% CIs. Infection and infestations include lung and urinary tract infections, COVID-19, and sepsis; gastrointestinal disorders include dysphagia, nausea, esophagitis; general disorders include fatigue, pain, and fever; respiratory, thoracic, and mediastinal disorders include dyspnea, cough, and pneumonitis; skin and subcutaneous tissue disorders include rash, dry skin, alopecia, and others. For further details refer to eTable 6 in Supplement 3. TRT indicates consolidative thoracic radiotherapy.

sure was not associated (HR, 0.57; 95% CI, 0.30-1.08; $P = .06$) with AE risk (eTable 10 in Supplement 3). Whether TRT was administered concurrently (ie, atezolizumab during the course of TRT) vs sequentially (ie, atezolizumab before or after TRT) was not associated with AE occurrence (eFigure 4 and eTable 11 in Supplement 3). While initial lymphocyte counts were similar in treatment arms, patients in arm A experienced persisting lymphocyte depletion after TRT, accompanied by an accumulation of AEs (**Table 2**; eFigure 4 in Supplement 3). This was specific for lymphocytes, while leukocyte and neutrophil counts were unaffected (eTable 12 in Supplement 3).

## Discussion

To our knowledge, these are the first results reported from a randomized phase 2 trial investigating TRT for ES-SCLC in the immunotherapy era. Addition of consolidative TRT to atezolizumab maintenance unexpectedly yielded a significantly higher incidence of partly fatal infection- and respiratory-related SAEs.

The primary end point was not met: median OS was numerically shorter in arm A vs B (6.7 vs 13.4 months; HR, 1.55; $P = .34$). Median OS and 1-year OS rate in arm B (13.4 months and 56.6%) exceeded the atezolizumab arm in IMpower133 (12.3 months and 51.7%), whereas arm A (6.7 months and 30.3%) fell below this benchmark.[1] The OS curves crossed at approximately 24 months, with 2-year rates numerically favoring arm A (24.2% vs 15.7%), confirmed in an updated OS analysis with extended follow-up, although this observation requires cautious interpretation given the small number of patients remaining at risk. A sensitivity analysis on an adjusted ITT excluding fatal AEs yielded consistent OS results, indicating that the unfavorable outcome in arm A is not solely associated with these fatal events.

PFS was similar between arms (2.4 vs 2.6 months; HR, 0.92; $P = .85$). This dissociation between PFS and OS is clinically informative: the absence of a PFS benefit argues against meaningful TRT-driven systemic tumor control, while the OS detriment in arm A (despite similar PFS) points to treatment-related toxic effects as primary driver for survival differences, rather than accelerated progression.

Prospective single-arm studies of TRT added to chemoimmunotherapy in ES-SCLC reported no increase in toxic effects.[6-8] However, in the randomized PACIFIC-2[12] and Checkmate-73L[13] trials, both investigating simultaneous tho-



Downloaded from jamanetwork.com by DKTZ-DT Krebsforschungszentrum user on 11/13/2026

Table 1. Baseline Patients Characteristics of the Intention-to-Treat Population, by Treatment Arm

| Characteristic | No. (%) | | |
|---|---|---|---|
| | Arm A (atezolizumab + TRT) | Arm B (atezolizumab only) | Total |
| Patients, total No. | 34 | 34 | 68 |
| Sex | | | |
| Female | 11 (32.4) | 14 (41.2) | 25 (36.8) |
| Male | 23 (67.6) | 20 (58.8) | 43 (63.2) |
| Age, mean (SD), y | 63.3 (9.1) | 65.6 (5.5) | 64.5 ( 7.5) |
| Smoking history | | | |
| Former | 22 (64.7) | 21 (61.8) | 43 (63.2) |
| Current | 11 (32.4) | 13 (38.2) | 24 (35.3) |
| Unknown | 1 (2.9) | 0 | 1 (1.5) |
| Thoracic response after induction therapy | | | |
| PR | 30 (88.2) | 26 (76.5) | 56 (82.4) |
| CR | 0 | 2 (5.9) | 2 (2.9) |
| Thoracic stable disease with extrathoracic PR | 3 (8.8) | 6 (17.6) | 9 (13.2) |
| Thoracic stable disease with extrathoracic CR | 1 (2.9) | 0 | 1 (1.5) |
| Prophylactic cranial irradiation | | | |
| No | 33 (97.1) | 33 (97.1) | 66 (97.1) |
| Yes | 1 (2.9) | 1 (2.9) | 2 (2.9) |
| Presence of brain metastases | | | |
| No | 27 (79.4) | 25 (73.5) | 52 (76.5) |
| Yes | 7 (20.6) | 9 (26.5) | 16 (23.5) |
| ECOG Performance Status | | | |
| 0 | 17 (50.0) | 10 (29.4) | 27 (39.7) |
| 1 | 17 (50.0) | 24 (70.6) | 41 (60.3) |
| Charlson Comorbidity Index | | | |
| <2 | 23 (67.6) | 22 (66.7) | 45 (67.2) |
| ≥2 | 11 (32.4) | 11 (33.3) | 22 (32.8) |
| Missing data, No. | 0 | 1 | 1 |
| **Pulmonary function, mean (SD), %** | | | |
| Vital capacity maximum | 81.9 (18.9) | 82.3 (18.0) | 82.1 (18.3) |
| FEV1 | 74.6 (12.5) | 69.5 (18.5) | 72.1 (15.8) |
| DLCO SB | 54.7 (12.7) | 53.1 (21.7) | 53.9 (17.7) |
| Missing data, No. | 1 | 1 | 2 |
| DLCO VA | 69.2 (16.0) | 67.5 (23.6) | 68.3 (20.0) |
| Missing data, No. | 1 | 1 | 2 |
| **Tumor size, mean (SD), mm** | | | |
| Lung | 33.1 (17.0) | 38.8 (30.1) | 36.0 (24.3) |
| Missing data, No.[a] | 12 | 12 | 24 |
| Mediastinal lymph nodes | 28.8 (14.0) | 28.8 (12.7) | 28.8 (13.1) |
| Missing data, No. | 19 | 16 | 35 |
| Overall tumor size in thorax | 48.0 (24.4) | 50.6 (34.2) | 49.4 (29.5) |
| Missing data, No. | 7 | 6 | 13 |

Abbreviations: CR, complete response; DLCO SB, diffusing capacity of the lung for carbon monoxide single breath; DLCO VA, diffusing capacity of the lung for carbon monoxide divided by the alveolar volume; ECOG, Eastern Cooperative Oncology Group; FEV1, forced expiratory volume in 1 second; PR, partial response; TRT, consolidative thoracic radiotherapy.

[a] Patients with no measurable thoracic tumor at baseline.

racic chemoradiotherapy with immunotherapy in non–small cell lung cancer (NSCLC), more fatal infections and toxic effects in the interventional arms were observed, consistent with our findings. Along these lines, TRT, but not a longer atezolizumab exposure in the absence of radiotherapy, was associated with increased AE risk in the TREASURE trial. The safety profile observed in arm B is largely consistent with published data on atezolizumab maintenance, whereas the rate of trAE grade 3 or greater (26%) in arm A clearly exceeds benchmarks for both atezolizumab monotherapy and consolidative TRT without immunotherapy (eTable 13 in Supplement 3), despite radiation doses and volumes within or below typical clini-



Downloaded from jamanetwork.com by DKTZ-DT Krebsforschungszentrum user on 11/13/2026

Table 2. Absolute Lymphocyte Count During the First 4 Cycles of Atezolizumab Maintenance in the Safety Population, by Treatment Arm

| Absolute lymphocyte count measure[a] | No. (%) | | | |
|---|---|---|---|---|
| | Arm A (atezolizumab + TRT) | Arm B (atezolizumab only) | Total | *P* value[b] |
| Patients, total No. | 31 | 33 | 64 | NA |
| Baseline | | | | |
| <0.001/μL | 8 (25.8) | 5 (15.2) | 13 (20.3) | .31 |
| Normal range | 22 (71.0) | 28 (84.8) | 50 (78.1) | |
| >0.048/μL | 1 (3.2) | 0 | 1 (1.6) | |
| **Atezolizumab maintenance cycle** | | | | |
| Cycle 1 | | | | |
| <0.001/μL | 7 (25.9) | 5 (20.0) | 12 (23.1) | .61 |
| Normal range | 20 (74.1) | 20 (80.0) | 40 (76.9) | |
| Cycle 2 | | | | |
| <0.001/μL | 22 (91.7) | 3 (13.6) | 25 (54.3) | <.001 |
| Normal range | 2 (8.3) | 18 (81.8) | 20 (43.5) | |
| >0.048/μL | 0 | 1 (4.5) | 1 (2.2) | |
| Cycle 3 | | | | |
| <0.001/μL | 14 (77.8) | 4 (26.7) | 18 (54.5) | .004 |
| Normal range | 3 (16.7) | 11 (73.3) | 14 (42.4) | |
| >0.048/μL | 1 (5.6) | 0 | 1 (3.0) | |
| Cycle 4 | | | | |
| <0.001/μL | 9 (69.2) | 1 (8.3) | 10 (40.0) | .002 |
| Normal range | 4 (30.8) | 11 (91.7) | 15 (60.0) | |

Abbreviations: NA, not applicable; TRT, thoracic radiotherapy (started after first administration of atezolizumab maintenance therapy).

[a] Normal ALC range, 0.001-0.048/μL (to convert to $\times 10^9$/L, multiply by 0.001).

[b] Per $\chi^2$ test.

cal ranges,[9,14-17] further supporting an interaction between the 2 modalities.

No significant differences in baseline characteristics distinguished patients with and without SAEs in arm A, nor did the concurrent vs sequential TRT timing. Among patients with fatal AEs, low baseline DLCO SB appeared as a potential risk factor, while other clinical and dosimetric parameters were not informative. However, these results should be interpreted cautiously given the small sample size. Furthermore, radiation-induced lymphopenia emerged as a potential risk factor for clinically impactful AEs. Pronounced and persistent lymphocyte depletion after TRT was accompanied by a higher incidence of AEs, predominantly infections and respiratory disorders. Taken together with the previous PACIFIC-2[12] and CheckMate-73L[13] results, these findings suggest that concurrent thoracic immunoradiotherapy may increase clinically relevant toxic effects without clear efficacy benefit, arguing against general use in both NSCLC and SCLC.

### Limitations

The main limitation of the study is the exploratory nature of analyses due to the premature termination. That being said, the unfavorable risk-benefit balance in the TREASURE trial affected both toxicity and efficacy outcomes, thereby retrospectively fully justifying the SMC's recommendation to halt the trial.

## Conclusions

In this randomized clinical trial, combining consolidative TRT with atezolizumab maintenance resulted in an unexpected increase of toxic effects, including partially fatal infections and pulmonary disorders, associated with a reduced baseline DLCO SB and protracted radiation-induced lymphopenia. Thus, in the maintenance setting after chemoimmunotherapy, addition of consolidative TRT cannot currently be recommended outside of clinical trials for unselected patients, and future studies exploring this strategy should consider mitigation strategies, such as dose exposure to highly perfused organs at risk (ie, lungs, heart, and large vessels), radiotherapy timing relative to immunotherapy cycles, and prospectively defined safety stopping rules. Ongoing biomarker analyses may assist in identifying subpopulations with particular risk or therapeutic benefit from this approach.

ARTICLE INFORMATION

**Accepted for Publication:** May 7, 2026.

**Published Online:** July 9, 2026. doi:10.1001/jamaoncol.2026.2330

**Open Access:** This is an open access article distributed under the terms of the CC-BY-NC-ND License, which does not permit alteration or commercial use, including those for text and data mining, AI training, and similar technologies. © 2026 Bozorgmehr F et al. *JAMA Oncology*.

**Author Affiliations:** Department of Thoracic Oncology, Thoraxklinik, Heidelberg University Hospital and National Center for Tumor Diseases (NCT), NCT Heidelberg, a partnership between German Cancer Research Center (DKFZ) and Heidelberg University Hospital, Heidelberg, Germany (Bozorgmehr, Chung, Thomas, Christopoulos); Translational Lung Research Center Heidelberg, German Center for Lung Research (DZL), Heidelberg, Germany (Bozorgmehr, Chung,



Downloaded from jamanetwork.com by DKTZ-DT Krebsforschungszentrum user on 11/13/2026

Heussel, Polke, Thomas, Christopoulos); Institute of Medical Biometry, Heidelberg University, Heidelberg, Germany (Behnisch, Merling); Department of Radiation Oncology, Heidelberg University Hospital, Heidelberg University (Weykamp); Heidelberg Institute of Radiation Oncology (HIRO), Heidelberg, Germany (Weykamp); National Center for Tumor Diseases (NCT), Heidelberg, Germany (Weykamp); Clinical Cooperation Unit Radiation Oncology, German Cancer Research Center (DKFZ), Heidelberg, Germany (Weykamp); Department of Radiation Oncology, University Hospital Marburg, Marburg, Germany (Adeberg); Department of Hematology and Oncology, University Medical Center Goettingen, Comprehensive Cancer Center Lower Saxony, Goettingen Site, Goettingen, Germany (Overbeck); Department of Radiotherapy and Radiation Oncology, University Medical Center Goettingen, Comprehensive Cancer Center Lower Saxony, Goettingen Site, Goettingen, Germany (El Shafie, Rieken); Department of Oncology and Hematology, Barmherzige Brüder Hospital Regensburg, Regensburg, Germany (Maguire); Department of Radiation Oncology, Barmherzige Brüder Hospital Regensburg, Regensburg, Germany (Koerber); Department of Thoracic Oncology, City of Cologne Municipal Hospitals, Lung Hospital Cologne Merheim, Cologne, Germany (Buchmeier); Department of Radiation Oncology, City of Cologne Municipal Hospitals, Cologne, Germany (Hammer-Hellmig); Department of Radiotherapy, University Hospital Essen, University of Duisburg-Essen, Essen, Germany (Gauler, Pöttgen); Department of Medicine I, University Hospital Carl Gustav Carus TU Dresden, Dresden, Germany, and National Center for Tumor Diseases, Dresden, Germany (Wermke); Department of Radiotherapy and Radiation Oncology, University Hospital Carl Gustav Carus TU Dresden, Dresden, Germany, and National Center for Tumor Diseases, Dresden, Germany (Troost); OncoRay–National Center for Radiation Research in Oncology, Faculty of Medicine and University Hospital Carl Gustav Carus TUD Dresden University of Technology, Dresden, Germany (Troost); Department of Pneumonology, Bogenhausen Hospital, Munich, Germany (Kokowski); Ambulatory Health Care Centre "Die Radiologie," Munich, Germany (Röper); Department of Thoracic Oncology and Palliative Care, DGD Lungenklinik Hemer, Hemer, Germany (Reinacher-Schick); Department of Hematology, Oncology and Palliative Care, St Josef-Hospital, Ruhr-University Bochum, Bochum, Germany (Reinacher-Schick); Department of Radiation Oncology, DGD Lungenklinik Hemer, Hemer, Germany (Consdorf); Department of Hematology/Oncology, RKH Hospital Ludwigsburg, Ludwigsburg, Germany (Ulmer); Department of Radiation Oncology and Radiation Therapy, RKH Hospital Ludwigsburg, Ludwigsburg, Germany (Müller); Department of Hematology/Oncology, Sana Klinikum Offenbach GmbH, Offenbach, Germany (Wehler); Department of Radiation Oncology, Sana Klinikum Offenbach GmbH, Offenbach, Germany (Hey-Koch); Department of Hematology, Medical Oncology, and Pneumology, University Medical Center Mainz, Mainz, Germany (Alt); Department of Radiation Oncology and Radiation Therapy, University Medical Center of the Johannes Gutenberg University Mainz, Mainz, Germany (Stockinger); Department of Thoracic Oncology, Lungenklinik Loewenstein, Loewenstein, Germany (Kuon); Department of Radiation Oncology, Klinikum Am Gesundbrunnen, SLK-Kliniken Heilbronn GmbH, Heilbronn, Germany (Bischof); Department of Pulmonary Medicine, Faculty of Medicine, Kepler University Hospital, Johannes-Kepler-University, Linz, Austria (Lamprecht); Department of Radiation Oncology, Ordensklinikum Linz GmbH, Barmherzige Schwestern, Linz, Austria (Geinitz); Joint Practice for Hematology and Oncology, Münster, Germany (Lerchenmüller); Medical Care Centre Strahlentherapie, Clemenshospital Münster, Münster, Germany (Kriz); Department of Pulmonology, Klinikum Klagenfurt Am Woerthersee, Klagenfurt, Austria (Rauter); Department of Radiation Oncology, Klinikum Klagenfurt, Klagenfurt, Austria (Raunik); Department of Internal Medicine, Pneumology, and Sleep Medicine, DRK Kliniken Berlin Mitte, Berlin, Germany (Schmidt); Department of Radiation Oncology, DRK Kliniken Berlin Westend, Berlin, Germany (Stupavsky); Department for Internal Medicine II, University of Ulm, Ulm, Germany (Kropf-Sanchen, Schmidtke-Schrezenmeier); Department of Radiotherapy and Radiation Oncology, University Hospital Ulm, Ulm, Germany (Wiegel); Department of Medicine V, University Hospital, Ludwig Maximilians University Munich, Munich, Germany (Tufman); Comprehensive Pneumology Center, German Center for Lung Research (DZL), Munich, Germany (Tufman); Department of Radiation Oncology, University Hospital Munich (LMU), Munich, Germany (Manapov); Diagnostic and Interventional Radiology, Heidelberg University Hospital, Heidelberg, Germany (Heussel); Department of Diagnostic and Interventional Radiology with Nuclear Medicine, Thoraxklinik, Heidelberg University Hospital, Heidelberg, Germany (Heussel); Center for Interstitial and Rare Lung Diseases, Department of Pneumology and Critical Care Medicine, Thoraxklinik, Heidelberg University Hospital, Heidelberg, Germany (Polke); Airway Research Center North, German Center for Lung Research, LungenClinic Grosshansdorf, Grosshansdorf, Germany (Reck); IKF, The Frankfurt Institute of Clinical Cancer Research, Frankfurt am Main, Germany (Al-Batran).

**Author Contributions:** Dr Bozorgmehr had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.
*Concept and design:* Bozorgmehr, Chung, Behnisch, Weykamp, El Shafie, Kuon, Tufman, Manapov, Al-Batran, Thomas, Christopoulos, Rieken.
*Acquisition, analysis, or interpretation of data:* Bozorgmehr, Chung, Behnisch, Adeberg, Overbeck, El Shafie, Maguire, Koerber, Buchmeier, Hammer-Hellmig, Gauler, Pöttgen, Wermke, Troost, Kokowski, Röper, Reinacher-Schick, Consdorf, Ulmer, Müller, Wehler, Hey-Koch, Alt, Stockinger, Bischof, Lamprecht, Geinitz, Lerchenmüller, Kriz, Rauter, Raunik, Schmidt, Stupavsky, Kropf-Sanchen, Schmidtke-Schrezenmeier, Wiegel, Tufman, Merling, Heussel, Polke, Reck, Al-Batran, Christopoulos, Rieken.
*Drafting of the manuscript:* Bozorgmehr, Chung, Behnisch, Weykamp, Pöttgen, Tufman, Manapov, Christopoulos, Rieken.
*Critical review of the manuscript for important intellectual content:* Bozorgmehr, Behnisch, Adeberg, Overbeck, El Shafie, Maguire, Koerber, Buchmeier, Hammer-Hellmig, Gauler, Wermke, Troost, Kokowski, Röper, Reinacher-Schick, Consdorf, Ulmer, Müller, Wehler, Hey-Koch, Alt, Stockinger, Kuon, Bischof, Lamprecht, Geinitz, Lerchenmüller, Kriz, Rauter, Raunik, Schmidt, Stupavsky, Kropf-Sanchen, Schmidtke-Schrezenmeier, Wiegel, Tufman, Manapov, Merling, Heussel, Polke, Reck, Al-Batran, Thomas, Christopoulos, Rieken.
*Statistical analysis:* Behnisch, Kokowski, Wehler, Merling, Christopoulos, Rieken.
*Obtained funding:* Bozorgmehr, Al-Batran, Thomas, Rieken.
*Administrative, technical, or material support:* Bozorgmehr, Chung, Adeberg, Koerber, Gauler, Pöttgen, Wermke, Troost, Kokowski, Reinacher-Schick, Kuon, Bischof, Kriz, Rauter, Heussel, Christopoulos, Rieken.
*Supervision:* Weykamp, Adeberg, Buchmeier, Ulmer, Müller, Lamprecht, Schmidtke-Schrezenmeier, Tufman, Reck, Al-Batran, Christopoulos, Rieken.
*Other:* Kropf-Sanchen.

**Conflict of Interest Disclosures:** Dr Bozorgmehr reported grants from Roche during the conduct of the study; grants from AstraZeneca; consulting fees from Immatics; honoraria from Novocure, Daiichi Sankyo, GlaxoSmithKline, and AstraZeneca; travel support from Janssen, Roche, and AstraZeneca; advisory board from Bristol Myers Squibb, AstraZeneca, Daiichi Sankyo, Janssen, Regeneron, and Pfizer outside the submitted work. Dr Behnisch reported a Thoraxklinik Cooperation agreement during the conduct of the study. Dr Weykamp reported personal fees from Varian Medical Systems, Merck Sharp & Dohme, AstraZeneca, and Novocure outside the submitted work. Dr Adeberg reported personal fees from AstraZeneca outside the submitted work. Dr Overbeck reported personal fees from commercial sponsor; honoraria for advisory boards from AstraZeneca, Bristol Myers Squibb, Boehringer Ingelheim, Daiichi Sankyo, Lilly, Janssen, Merck KGaA, Merck Sharp & Dohme, Novartis, Pfizer, Pierre Fabre, Roche, and Takeda and personal fees from commercial sponsor; travel support from Amgen, AstraZeneca, Boehringer Ingelheim, Daiichi Sankyo, and Janssen outside the submitted work. Dr El Shafie reported grants from AstraZeneca during the conduct of the study; personal fees from AstraZeneca, Novocure, Merck, Merck Sharp & Dohme, and personal fees from Accuray outside the submitted work. Dr Koerber reported personal fees from Think Wired! and nonfinancial support from AstraZeneca outside the submitted work. Dr Buchmeier reported personal fees from AstraZeneca, Bristol Myers Squibb, Johnson & Johnson, Regeneron, and Roche outside the submitted work. Dr Gauler reported personal fees from Roche AG, Merck Sharp & Dohme, and AstraZeneca during the conduct of the study; personal fees from Eisai and Ipsen outside the submitted work. Dr Pöttgen reported honoraria from AstraZeneca outside the submitted work. Dr Wermke reported personal fees from Lilly and Boehringer Ingelheim; grants from Boehringer Ingelheim; personal fees from SYNLAB, Merck Serona, Amgen, Novartis, Pfizer, Bristol Myers Squibb, Regeneron, MJH/PER, AstraZeneca, Takeda, and Daiichi Sankyo; grants from Roche and Iovance; personal fees from Iovance, ISA Pharmaceuticals, Novartis, PharmaMar, Tknife, Tacalyx, Zymeworks, Immatics, and Genentech outside the submitted work. Dr Troost reported



Downloaded from jamanetwork.com by DKTZ-DT Krebsforschungszentrum user on 11/13/2026

personal fees from IBA for being a member of SAB outside the submitted work. Dr Reinacher-Schick reported personal fees from Amgen, Pierre Fabre, Roche, Takeda, AstraZeneca, Hoffmann-La Roche, and MCI during the conduct of the study; grants from Roche, AIO Studien, ARCUS Biosciences Europe Dublin, BioNTech, Charité Universitätsmedizin Berlin, Genentech, Georg-August-Universität Göttingen, Johannes-Gutenberg-Universität Mainz, Merck Healthcare Darmstadt, Rafael Pharmaceutics, Resilience Paris, and Universitätsklinikum Ulm outside the submitted work. Dr Ulmer reported presentations for Roche outside the submitted work. Dr Müller reported nonfinancial support from AstraZeneca, Medea, and Vision RT outside the submitted work. Dr Wehler reported institutional fees from Frankfurt Institute of Clinical Cancer Research during the conduct of the study and grants from Frankfurt Institute of Clinical Cancer Research outside the submitted work. Dr Alt reported personal fees from Roche during the conduct of the study. Dr Geinitz reported grants from AstraZeneca outside the submitted work. Dr Kropf-Sanchen reported honoraria and advisory role from Amgen, AstraZeneca, BeOne, Boehringer Ingelheim, Bristol Myers Squibb, Daiichi Sankyo, Johnson & Johnson, Merck Sharp & Dohme, Novartis, Pfizer, Regeneron, Roche, Takeda, Onkowissen.TV, Art Tempi/Medtoday, and Streamed Up; travel support from AstraZeneca, Amgen, Boehringer Ingelheim, Novartis, Roche, Pfizer, Johnson & Johnson; grants from Hector Foundation II; serving as chairman of the pneumological oncology working group of the German Cancer Society, chairman of the AIO working group on women and family support in Oncology of the German Cancer Society, member of the thoracic oncology steering committee of the internal oncology working group of the German Cancer Society, managing director of the south German society for pneumology. Dr Tufman reported grants from AstraZeneca outside the submitted work; and consulting and speakers fees for Lilly, Roche, Novartis, Pfizer, Amgen, Merck Sharp & Dohme, Bristol Myers Squibb, Takeda, BeOne, Regeneron, AstraZeneca, and GlaxoSmithKline outside the submitted work. Ms Merling reported a cooperation agreement from Thoraxklinik during the conduct of the study. Dr Heussel reported personal fees from Boehringer-Ingelheim, Voiant, Exscientia, AstraZeneca, Pfizer, Chiesi, and Sanofi outside the submitted work. Dr Polke reported consulting/speaking/travel fees from Apontis Pharma, AstraZeneca, Boehringer Ingelheim, and Novartis outside the submitted work. Dr Al-Batran reported grants from Roche during the conduct of the study; equity in Frankfurt Institute of Clinical Cancer Research; grants from Merck Sharp & Dohme, Eli Lilly Germany, Bristol Myers Squibb, MCI Deutschland, Celgene, Vifor Pharma, Sanofi, German Cancer Aid (Krebshilfe), German Research Foundation, Federal Ministry of Education and Research of Germany, Roche, Eurozyto, Immutep, Ipsen, AstraZeneca, Merck Sharp & Dohme, Eli Lilly Germany, and Bristol Myers Squibb outside the submitted work. Dr Thomas reported grants from Roche during the conduct of the study and outside the submitted work. Dr Christopoulos reported grants from Roche during the conduct of the study; grants from Takeda, AstraZeneca, Amgen, Johnson & Johnson, PharmaMar, Novartis, Boehringer Ingelheim, Roche, AstraZeneca, Roche, Takeda, Johnson & Johnson, Pfizer, BeOne, Takeda, MSD, Boehringer Ingelheim, Novartis, Daiichi Sankyo, Gilead, and ThermoFisher outside the submitted work. Dr Rieken reported grants from Stiftung Deutsche Krebshilfe, Bundesamt für Strahlenschutz, Land Niedersachsen, Volkswagen Stiftung, AstraZeneca, Novocure, Med update, Onkowissen, Lilly, and Roche outside the submitted work. No other disclosures were reported.

**Funding/Support:** The TREASURE trial was sponsored by the IKF, Frankfurt Institute of Clinical Cancer Research, and funded by a research grant from Roche.

**Role of the Funder/Sponsor:** The funder (Roche) had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication. The sponsor (IKF Frankfurt) contributed to preparation of the study protocol, study management and collection and management of the data. This trial was conducted by the Thoracic Oncology Working Group of the Arbeitsgemeinschaft Internistische Onkologie within the German Cancer Society (AIO-TRK-0320).

**Data Sharing Statement:** See Supplement 4.

**Additional Contributions:** The authors thank all the participants enrolled in this study and their families and caregivers. The authors also thank the physicians and nurses who cared for the participants and the staff at the clinical sites.

## REFERENCES

1. Horn L, Mansfield AS, Szczęsna A, et al; IMpower133 Study Group. First-line atezolizumab plus chemotherapy in extensive-stage small-cell lung cancer. *N Engl J Med*. 2018;379(23):2220-2229. doi:10.1056/NEJMoa1809064

2. De Ruysscher D, Reynders K, Van Limbergen E, Lambrecht M. Radiotherapy in combination with immune checkpoint inhibitors. *Curr Opin Oncol*. 2017;29(2):105-111. doi:10.1097/CCO.0000000000000352

3. Lin SH, Lin Y, Yao L, et al. Phase II trial of concurrent atezolizumab with chemoradiation for unresectable NSCLC. *J Thorac Oncol*. 2020;15(2):248-257. doi:10.1016/j.jtho.2019.10.024

4. Welsh J, Menon H, Chen D, et al. Pembrolizumab with or without radiation therapy for metastatic non-small cell lung cancer: a randomized phase I/II trial. *J Immunother Cancer*. 2020;8(2):e001001. doi:10.1136/jitc-2020-001001

5. Bozorgmehr F, Chung I, Fischer JR, et al. Reconsidering palliative radiotherapy in addition to PD-1 blockade for non-small cell lung cancer: results from the FORCE phase II trial (AIO/YMO-TRK-0415). *Clin Exp Metastasis*. 2025;42(5):42. doi:10.1007/s10585-025-10358-x

6. Welsh JW, Heymach JV, Chen D, et al. Phase I trial of pembrolizumab and radiation therapy after induction chemotherapy for extensive-stage small cell lung cancer. *J Thorac Oncol*. 2020;15(2):266-273. doi:10.1016/j.jtho.2019.10.001

7. Perez BA, Kim S, Wang M, et al. Prospective single-arm phase 1 and 2 study: ipilimumab and nivolumab with thoracic radiation therapy after platinum chemotherapy in extensive-stage small cell lung cancer. *Int J Radiat Oncol Biol Phys*. 2021;109(2):425-435. doi:10.1016/j.ijrobp.2020.09.031

8. Addeo A, Dietrich D, Mach N, et al. 1791P Thoracic radiotherapy plus maintenance durvalumab after first-line carboplatin and etoposide plus durvalumab in extensive-stage disease small cell lung cancer (ED-SCLC): A multicenter single-arm open-label phase II trial (SAKK 15/19)tle. *Ann Oncol*. 2024;35:S1065. doi:10.1016/j.annonc.2024.08.1882

9. Slotman BJ, van Tinteren H, Praag JO, et al. Use of thoracic radiotherapy for extensive stage small-cell lung cancer: a phase 3 randomised controlled trial. *Lancet*. 2015;385(9962):36-42. doi:10.1016/S0140-6736(14)61085-0

10. Bozorgmehr F, Christopoulos P, Chung I, et al. Protocol of the TREASURE study: thoracic radiotherapy with atezolizumab in small cell lung cancer extensive disease: a randomized, open-label, multicenter phase II trial. *BMC Cancer*. 2022;22(1):1011. doi:10.1186/s12885-022-10074-9

11. Bozorgmehr F, Weykamp F, Overbeck TR, et al. 1988MO Recruitment discontinuation in TREASURE trial (thoracic radiotherapy with atezolizumab in small cell lung cancer extensive disease) due to unexpected safety data. *Ann Oncol*. 2023;34:S1060. doi:10.1016/j.annonc.2023.09.1219

12. Bradley JD, Sugawara S, Lee KH, et al; PACIFIC-2 Investigators. Simultaneous durvalumab and platinum-based chemoradiotherapy in unresectable stage iii non-small cell lung cancer: the phase III PACIFIC-2 study. *J Clin Oncol*. 2025;43(33):3610-3621. doi:10.1200/JCO-25-00036

13. Peters S, Tan D, Gerber DE, et al. 650 CheckMate 73L: Phase III study comparing nivolumab (N) + concurrent chemoradiotherapy (CCRT) followed by N ± ipilimumab (I) v CCRT followed by durvalumab (D) for previously untreated, locally advanced stage (stg) III NSCLC. *Immunooncol Technol*. 2024;24:100808. doi:10.1016/j.iotech.2024.100808

14. Yu J, Jiang L, Zhao L, et al; Trial Management Group. High-dose hyperfractionated simultaneous integrated boost radiotherapy versus standard-dose radiotherapy for limited-stage small-cell lung cancer in China: a multicentre, open-label, randomised, phase 3 trial. *Lancet Respir Med*. 2024;12(10):799-809. doi:10.1016/S2213-2600(24)00189-9

15. Salama JK, Pang H, Bogart JA, et al. Predictors of pulmonary toxicity in limited stage small cell lung cancer patients treated with induction chemotherapy followed by concurrent platinum-based chemotherapy and 70 Gy daily radiotherapy: CALGB 30904. *Lung Cancer*. 2013;82(3):436-440. doi:10.1016/j.lungcan.2013.10.001

16. Grønberg BH, Killingberg KT, Fløtten Ø, et al. High-dose versus standard-dose twice-daily thoracic radiotherapy in limited-stage SCLC: final survival data, long-term toxicity, and relapse patterns in a randomized, open-label, phase II trial. *J Thorac Oncol*. 2025;20(8):1108-1119. doi:10.1016/j.jtho.2025.04.007

17. Kandler C, Elsayad K, Evers G, et al. Reduction of tumor volume during radiotherapy in patients with small-cell lung cancer and its prognostic significance. *Strahlenther Onkol*. 2023;199(11):1011-1017. doi:10.1007/s00066-023-02146-x



Downloaded from jamanetwork.com by DKTZ-DT Krebsforschungszentrum user on 11/13/2026